

## 8-hour time-restricted eating linked to a 91% higher risk of cardiovascular death

March 18 2024



Credit: Unsplash/CC0 Public Domain



An analysis of over 20,000 U.S. adults found that people who limited their eating to less than 8 hours per day, a time-restricted eating plan, were more likely to die from cardiovascular disease compared to people who ate across 12-16 hours per day, according to preliminary research presented at the American Heart Association's Epidemiology and Prevention | Lifestyle and Cardiometabolic Scientific Sessions 2024, March 18- 21, in Chicago. The meeting offers the latest science on population-based health and wellness and implications for lifestyle.

Time-restricted eating, a type of intermittent fasting, involves limiting the hours for eating to a specific number of hours each day, which may range from a 4- to 12-hour time window in 24 hours. Many people who follow a time-restricted eating diet follow a 16:8 eating schedule, where they eat all their foods in an 8-hour window and fast for the remaining 16 hours each day, the researchers noted.

Previous research has found that time-restricted eating improves several cardiometabolic health measures, such as blood pressure, blood glucose, and cholesterol levels.

"Restricting daily eating time to a short period, such as 8 hours per day, has gained popularity in recent years as a way to lose weight and improve heart health," said senior study author Victor Wenze Zhong, Ph.D., a professor and chair of the department of epidemiology and biostatistics at the Shanghai Jiao Tong University School of Medicine in Shanghai, China.

"However, the long-term health effects of time-restricted eating, including the risk of death from any cause or <u>cardiovascular disease</u>, are unknown."



In this study, researchers investigated the potential long-term health impact of following an 8-hour time-restricted eating plan. They reviewed information about <u>dietary patterns</u> for participants in the annual 2003-2018 National Health and Nutrition Examination Surveys (NHANES) in comparison to data about people who died in the U.S. from 2003 through December 2019 from the Centers for Disease Control and Prevention's National Death Index database.

The analysis found:

- People who followed a pattern of eating all of their food across less than 8 hours per day had a 91% higher risk of death due to cardiovascular disease.
- The increased risk of cardiovascular death was also seen in people living with heart disease or cancer.
- Among people with existing cardiovascular disease, an eating duration of no less than 8 but less than 10 hours per day was also associated with a 66% higher risk of death from heart disease or stroke.
- Time-restricted eating did not reduce the overall risk of death from any cause.
- An eating duration of more than 16 hours per day was associated with a lower risk of cancer mortality among people with cancer.

"We were surprised to find that people who followed an 8-hour, timerestricted eating schedule were more likely to die from cardiovascular disease. Even though this type of diet has been popular due to its potential short-term benefits, our research clearly shows that, compared with a typical eating time range of 12-16 hours per day, a shorter eating duration was not associated with living longer," Zhong said.

"It's crucial for patients, particularly those with existing heart conditions or cancer, to be aware of the association between an 8-hour eating



window and increased risk of cardiovascular death. Our study's findings encourage a more cautious, personalized approach to dietary recommendations, ensuring that they are aligned with an individual's health status and the latest scientific evidence," he continued.

"Although the study identified an association between an 8-hour eating window and cardiovascular death, this does not mean that time-restricted eating caused cardiovascular death."

Study details and background:

- The study included approximately 20,000 adults in the U.S. with an average age of 49 years.
- Study participants were followed for a median length of 8 years and a maximum length of 17 years.
- The study included data for NHANES participants who were at least 20 years old at enrollment between 2003-2018 and had completed two 24-hour dietary recall questionnaires within the first year of enrollment.
- Approximately half of the participants self-identified as men, and half self-identified as women. 73.3% of the participants selfidentified as non-Hispanic white adults, 11% self-identified as Hispanic adults, 8% self-identified as non-Hispanic Black adults, and 6.9% of adults self-identified as another racial category, including mixed-race adults and adults of other non-Hispanic races.

The study's limitations included its reliance on self-reported dietary information, which may be affected by participant's memory or recall and may not accurately assess typical eating patterns. Factors that may also play a role in health, outside of the daily duration of eating and cause of death, were not included in the analysis.



Future research may examine the biological mechanisms that underly the associations between a time-restricted eating schedule and adverse cardiovascular outcomes, and whether these findings are similar for people who live in other parts of the world, the authors noted.

"Overall, this study suggests that time-restricted eating may have shortterm benefits but long-term adverse effects. When the study is presented in its entirety, it will be interesting and helpful to learn more of the details of the analysis," said Christopher D. Gardner, Ph.D., FAHA, the Rehnborg Farquhar Professor of Medicine at Stanford University in Stanford, California, and chair of the writing committee for the Association's 2023 scientific statement, <u>Popular Dietary Patterns:</u> <u>Alignment with American Heart Association 2021 Dietary Guidance</u>.

"One of those details involves the nutrient quality of the diets typical of the different subsets of participants. Without this information, it cannot be determined if nutrient density might be an alternate explanation to the findings that currently focus on the window of time for eating. Second, it needs to be emphasized that categorization into the different windows of time-restricted eating was determined on the basis of just two days of dietary intake," he said.

"It will also be critical to see a comparison of demographics and baseline characteristics across the groups that were classified into the different time-restricted eating windows—for example, was the group with the shortest time-restricted eating window unique compared to people who followed other eating schedules, in terms of weight, stress, traditional cardiometabolic risk factors or other factors associated with adverse cardiovascular outcomes?"

"This additional information will help to understand better the potential independent contribution of the short time-restricted eating pattern reported in this interesting and provocative abstract."



## Provided by American Heart Association

Citation: 8-hour time-restricted eating linked to a 91% higher risk of cardiovascular death (2024, March 18) retrieved 9 May 2024 from <u>https://medicalxpress.com/news/2024-03-hour-restricted-linked-higher-cardiovascular.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.