

New imaging technology for diagnosing and treating bowel cancer patients

March 19 2024



Representative transverse PET images from each mouse model and tracer. Credit: *Clinical Cancer Research* (2024). DOI: 10.1158/1078-0432.CCR-23-1063

Biopsies are currently used to diagnose bowel cancer but require an



invasive procedure that offers risks, such as potential infection, and are unable to capture an entire picture of what is happening in a patient's bowel.

New research from the Cancer Research UK Scotland Institute and researchers at the University of Glasgow, <u>published</u> in *Clinical Cancer Research*, found that positron emission tomography (PET) imaging could allow the entire bowel to be examined and tumors studied while inside the body, rather than examining tumor tissue once it has been removed.

PET scanning creates a three-dimensional picture of the inside of the body using labeled nutrients, or drugs known as a tracer, to show up areas where cells are more active than normal.

The technology would allow bowel cancer patients to be matched to the best treatment for them, which is an area known as precision medicine, a growing area of oncology.

The team also believes multiple PET scans during a patient's treatment could enable the progression of the cancer, and impact of treatment, to be effectively monitored.

Dr. David Lewis, of the Cancer Research UK Scotland Institute and the University of Glasgow, who led the research, said, "Precision medicine has the potential to revolutionize <u>cancer diagnosis</u> and treatment. However, the development of accurate, informative, and patient-friendly diagnostic techniques is crucial for its success.

"PET imaging offers a promising alternative, with the ability to survey the entire cancer landscape, allowing us to examine tumors in more detail while they are still growing."

Each year about 4,000 people are diagnosed with bowel cancer in



Scotland. With about 1,800 Scots sadly losing their lives to the disease annually, the search for new and better treatments is vital.

The research team used existing <u>genetic data</u> on bowel cancer to identify different tumor characteristics using PET imaging.

They also discovered that multiple tracer PET scanning, which uses different PET tracers, rather than just one, was able to distinguish between different types of bowel cancer in mice based on their genes.

This would allow a more personalized approach for patients as treatments could be tailored to their particular cancer.

Patients can have different mutations in their bowel cancer, and mutations in genes such as KRAS, APC and TGFB have different image signatures.

The team believes PET imaging could identify the type of bowel cancer a patient has, by matching the image signature with the mutations, therefore allowing the patient to access the best treatment for their individual disease more quickly.

Dr. Catherine Elliott, Director of Research at Cancer Research UK, said, "These findings by the team at the Cancer Research UK Scotland Institute and University of Glasgow offer an exciting opportunity to revolutionize the way we diagnose and monitor <u>bowel cancer</u> without invasive surgery, reducing the risk and improving outcomes for patients.

"PET imaging is a crucial tool in our future approach to diagnosing this disease which affects so many people in Scotland."



More information: Gaurav Malviya et al, Noninvasive Stratification of Colon Cancer by Multiplex PET Imaging, *Clinical Cancer Research* (2024). DOI: 10.1158/1078-0432.CCR-23-1063

Provided by University of Glasgow

Citation: New imaging technology for diagnosing and treating bowel cancer patients (2024, March 19) retrieved 8 May 2024 from <u>https://medicalxpress.com/news/2024-03-imaging-technology-bowel-cancer-patients.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.