

# Data show even low levels of leisure time physical activity help to lower stroke risk

March 5 2024

---



Credit: CC0 Public Domain

Even people whose physical activity levels fall short of recommended guidelines, but who manage to do some during their leisure time, are likely to have a lower risk of stroke than their sedentary peers, suggests a

pooled data analysis of the available evidence, published online in the *Journal of Neurology, Neurosurgery & Psychiatry*.

The effects are independent of age and sex, the findings show, prompting the authors to suggest that everyone should be encouraged to do whatever level of [physical activity](#) they can manage in their [leisure time](#).

There's no doubt that moderate to high levels of leisure-time physical activity curb stroke risk. But it's not clear if even modest amounts confer protection, and if any such effects depend on age and sex, explain the authors.

While international guidelines recommend 150 minutes or more a week of moderate intensity physical activity or 75 minutes or more of vigorous intensity activity to lower the risk of cardiovascular disease, including stroke, not many adults achieve this, say the researchers.

To find out if lower levels of physical activity might still confer protection, the authors trawled research databases for relevant studies. They pooled the results of 15, involving 752,050 adults whose health had been monitored for an average of 10.5 years.

Each study assessed the potential impact of between three—none, below target, and ideal—and five—none, insufficient, low, moderate and intense— levels of leisure time physical activity on stroke risk.

The pooled data analysis of five studies assessing three levels of leisure time physical activity showed that, compared with no physical activity, the highest 'ideal' amount cut the risk of stroke by 29%, but that some 'below target' activity still reduced the risk by 18%.

Similar findings emerged for the pooled data analysis of the six studies

reporting on four, and the two reporting on five levels of leisure time physical activity. Compared with none, a moderate level of physical activity cut the risk of stroke by between 27% and 29%.

The authors acknowledge several limitations to their findings, chief among which were the variable definitions of different levels of activity used in the included studies and the reliance on subjectively assessed levels of physical activity.

Nevertheless, the authors conclude that recreational physical activity, even in small amounts could help ward off stroke in the long term.

"According to our results, all levels of [leisure time physical activity] can be beneficial for stroke prevention, including levels currently regarded as low or insufficient," they write. "People should be encouraged to be physically active even at the lowest levels."

**More information:** Risk of stroke with different levels of leisure-time physical activity: a systematic review and meta-analysis of prospective cohort studies, *Journal of Neurology, Neurosurgery & Psychiatry* (2024). [DOI: 10.1136/jnnp-2023-332457](https://doi.org/10.1136/jnnp-2023-332457)

Provided by British Medical Journal

Citation: Data show even low levels of leisure time physical activity help to lower stroke risk (2024, March 5) retrieved 29 April 2024 from <https://medicalxpress.com/news/2024-03-leisure-physical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.