

Running performance helped by mathematical research

March 5 2024



Simulated running speed for Jakob Ingebrigtsen: during his race in Munich (blue line), with the maximal value of VO₂ increased by 5% (green line), with anaerobic energy e^0 increased by 5% (red line), with the maximal propulsive force f^M increased by 10% (black line), and with kinetics increased by 5% (purple line). Credit: Amandine Aftalion

How to optimize running? A new mathematical model has shown, with great precision, the impact that physiological and psychological parameters have on running performance and provides tips for optimized



training. The model grew out of research conducted by a French-British team including two CNRS researchers, the <u>results</u> of which appear in the journal *Frontiers in Sports and Active Living*.

This innovative model was developed thanks to extremely precise data from the performances of Matthew Hudson-Smith (400m), Femke Bol (400m), and Jakob Ingebrigtsen (1500m) at the 2022 European Athletics Championships in Munich, and for Gaia Sabbatini (1500m) at the 2021 European Athletics U23 Championships in Tallinn. It led to an optimal control problem for finishing time, effort, and energy expenditure.

This is the first time that such a model has also considered the variability of motor control, i.e., the role of the brain in the process of producing movement. The simulations allow the researchers to have access to the physiological parameters of the runners—especially <u>oxygen consumption</u> (or VO₂), and <u>energy expenditure</u> during the race—as well as compute their variations. Quantifying costs and benefits in the model provides immediate access to the best strategy for achieving the runner's optimal performance.

The study details multiple criteria, such as the importance of a quick start in the first 50 meters (due to the need for fast oxygen kinetics), or reducing the decrease in velocity in a 400m race. The scientists also demonstrated that improving the aerobic metabolism (oxygen uptake) and the ability to maintain VO₂ are crucial elements to 1500m race performance.

The development of this model represents considerable progress in studying variations in physiological parameters during championship races, for which in vivo measurements are not possible.

More information: Antoine Le Hyaric et al, Modelling the optimization of world-class 400 m and 1,500 m running performances



using high-resolution data, *Frontiers in Sports and Active Living* (2024). DOI: 10.3389/fspor.2024.1293145

Provided by CNRS

Citation: Running performance helped by mathematical research (2024, March 5) retrieved 13 May 2024 from <u>https://medicalxpress.com/news/2024-03-mathematical.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.