


# New studies suggest millions with mild cognitive impairment go undiagnosed, often until it's too late

March 24 2024, by Soeren Mattke and Ying Liu

---



Credit: Pixabay/CC0 Public Domain

Mild cognitive impairment—an early stage of dementia—is widely

underdiagnosed in people 65 and older. That is the key takeaway of two recent studies from our team.

In the first study, we used Medicare data for about 40 million beneficiaries age 65 and older from 2015 to 2019 to estimate the prevalence of mild cognitive impairment in that population and to identify what proportion of them had actually been diagnosed.

Our [finding was sobering](#): A mere 8% of the number of cases with mild cognitive impairment that we expected based on a statistical model had actually been diagnosed. Scaled up to the general population 65 and older, this means that approximately 7.4 million cases across the country remain undiagnosed.

In the second study, we analyzed data for 226,756 primary care clinicians and found that [over 99% of them underdiagnosed mild cognitive impairment](#) in this population.

## Why it matters

Mild cognitive impairment is an early symptom of Alzheimer's disease in [about half of cases](#) and progresses to dementia [at a rate of 10% to 15% per year](#). It includes symptoms such as losing the ability to remember recent events and appointments, make sound decisions and master complex tasks. Failure to detect it might deprive patients of an opportunity to get treated and to slow down [disease progression](#).

Mild cognitive impairment can sometimes be caused by easily addressable factors, such as medication side effects, thyroid dysfunction or vitamin B12 deficiency. Since mild cognitive impairment has [the same risk factors as cardiovascular disease](#), such as [high blood pressure](#) and cholesterol, medication management of these risks combined with diet and exercise [can reduce the risk of progression](#).

In 2023, the Food and Drug Administration [approved the drug lecanemab](#) as the first disease-modifying treatment [for Alzheimer's disease](#), the most common cause of mild cognitive impairment. In contrast to previous drugs, which can temporarily improve symptoms of the disease, such as [memory loss](#) and agitation, this new treatment addresses the underlying cause of the disease.

Lecanemab, a monoclonal antibody, [reduces amyloid plaques](#) in the brain, which are toxic protein clumps that are believed to contribute to the progression of the disease. In a large clinical trial, lecanemab was able to [reduce the progression](#) of early-stage Alzheimer's disease. A similar drug, donanemab, also [succeeded in a clinical trial](#) and is expected to be [approved sometime in 2024](#).

However, these drugs must be used in the early stages of Alzheimer's disease, ideally when a patient has only mild cognitive impairment, as there is [no evidence that they are effective in advanced stages](#).

## What still isn't known

Many factors contribute to the [lack of timely detection](#). But researchers don't have a good understanding of the relative importance of those individual factors or how to reduce the high rate of underdiagnosis.

While distinct, symptoms are subtle and their slow progression means that they can be overlooked or misinterpreted as normal aging. A neurologist in China told our research team that diagnosis rates spike in China after the New Year's holiday, when children who haven't seen their parents for a year notice changes that are harder to pick up when interacting with someone daily.

Doctors also commonly discount memory concerns as normal aging and doubt that much can be done about it. While cognitive tests to

distinguish [mild cognitive impairment](#) from pathologic decline do exist, they take about 15 minutes, which can be hard to come by during the limited time of a doctor's visit and may require a follow-up appointment.

## What's next

People, particularly those in their 60s and beyond, as well as their families and friends need to be vigilant about cognitive decline, bring it up during doctor's appointments and insist on a formal assessment.

The [Medicare yearly "wellness" visit](#) is an opportunity to explore such concerns, but only about half of beneficiaries [take advantage of it](#).

Just as physicians ask patients about unexplained weight loss and take those concerns seriously, we believe questions that explore a patient's cognitive state need to become the norm.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: New studies suggest millions with mild cognitive impairment go undiagnosed, often until it's too late (2024, March 24) retrieved 27 April 2024 from <https://medicalxpress.com/news/2024-03-millions-mild-cognitive-impairment-undiagnosed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.