

# Patient recovery after surgery for esophageal cancer isn't influenced by using standard or keyhole incisions

March 25 2024

---



Credit: Pixabay/CC0 Public Domain

New research has found no evidence of a difference between recovery time and complications when comparing standard and keyhole surgical incisions for the treatment of oesophageal cancer (cancer of the gullet). The [study](#), led by the University of Bristol Medical School and published in the *British Journal of Surgery*, showed surgeons treating patients with oesophageal cancer do not need to change their practice if they have a strong preference for either procedure type.

Oesophageal [cancer](#) is the tenth most common cancer globally. It causes one in 18 cancer-related deaths. If only the esophagus and local lymph nodes are affected, surgeons usually perform a procedure called an oesophagectomy. This means they remove the esophagus to try and cure the cancer. The two most common ways of doing this use either standard incisions (two large cuts) or 'keyhole' incisions (one large cut and several small ones)

The ROMIO study involved patients being randomly assigned to two groups. One of the groups had standard surgery (263 people), and the other had keyhole surgery (264 people).

Researchers found no differences between the groups in relation to:

- recovery three months after surgery, as measured by patient-completed questionnaires about physical function
- how often patients developed complications and how severe the complications were
- the extent to which the cancer was removed (it was equally well removed in both groups)

Chris Metcalfe, Professor of Medical Statistics at Bristol Medical School: Population Health Sciences (PHS), said, "Our study didn't

confirm findings from previous trials, which suggested that minimally invasive approaches to oesophagectomy reduced the number of complications patients would develop."

"We found no evidence of differences between standard and keyhole approaches in relation to short-term clinical outcomes or patient-reported recovery of physical function over three months. There was no strong evidence that the costs of NHS resources in the first three months differed between the two approaches to the procedure."

"We will report about the longer-term recovery (24-month follow-up) and health of ROMIO participants in a separate publication. We will also publish the findings from a nested study on totally keyhole surgery."

Jane Blazeby, Professor of Surgery in the Bristol Medical School: PHS, added, "These results show us that recovery after this type of major surgery is not influenced by the type of incision used by surgeons, despite prior beliefs that the keyhole approach is better than standard incisions.

"Future research is now needed to carefully evaluate robotic surgical techniques for oesophageal cancer surgery."

**More information:** Chris Metcalfe et al, Laparoscopic or open abdominal surgery with thoracotomy for patients with oesophageal cancer: ROMIO randomized clinical trial, *British Journal of Surgery* (2024). [DOI: 10.1093/bjs/znae023](https://doi.org/10.1093/bjs/znae023)

Provided by University of Bristol

Citation: Patient recovery after surgery for esophageal cancer isn't influenced by using standard

or keyhole incisions (2024, March 25) retrieved 8 May 2024 from  
<https://medicalxpress.com/news/2024-03-patient-recovery-surgery-esophageal-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.