


Patients with delirium more likely to develop dementia, finds study



Association of delirium with death by baseline group (upper panel) and episodes of delirium recorded within first 12 months of follow-up (landmark period; lower panel). Credit: *BMJ* (2024). DOI: 10.1136/bmj-2023-077634

University of Queensland researchers have found older patients who experience delirium are three times more likely to develop dementia. Professor Ruth Hubbard from UQ's Center for Health Services Research (CHSR) said the study also found that every episode of delirium increased the chance of developing dementia by 20%.

March 28 2024



"Delirium is an acute confusional state which particularly impacts older people and can have long term detrimental effects on patients," Professor Hubbard said. "Our study found that it is also strongly linked to <u>dementia</u>."

The researchers analyzed a cohort of more than 110,000 patients from New South Wales hospitals over five years. The <u>research paper</u> was published in the *British Medical Journal*.

Lead author and CHSR research fellow, Dr. Emily Gordon, said more than 55,000 patients who had experienced delirium were compared to patients who had not.

"We paired the patients by matching certain criteria such as age, sex, frailty, reason for being in hospital, length of stay in hospital and length of stay in the <u>intensive care unit</u>," Dr. Gordon said.

"This was done to ensure we got as close as possible to isolating the impact of delirium on dementia risk, and to ignore the effects of other factors known to impact dementia, such as older age. We then followed both cohorts over five years to see whether they were eventually diagnosed with dementia."

Dr. Gordon said the findings highlighted the need for delirium prevention and care. "Up to 40% of delirium cases are preventable, and treatments are readily available. Measures to prevent delirium include keeping patients well nourished, hydrated and mobile.

"This study highlights the importance of delirium care and prevention—if we can reduce <u>delirium</u>, we may be able to reduce the global burden of dementia."

The data for this study was managed by research partners at the



Australian Institute of Health Innovation at Macquarie University.

More information: Emily H Gordon et al, Delirium and incident dementia in hospital patients in New South Wales, Australia: retrospective cohort study, *BMJ* (2024). DOI: 10.1136/bmj-2023-077634

Provided by University of Queensland

Citation: Patients with delirium more likely to develop dementia, finds study (2024, March 28) retrieved 8 May 2024 from https://medicalxpress.com/news/2024-03-patients-delirium-dementia.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.