


Study: PR55 α -controlled PP2A inhibits p16 expression and blocks cellular senescence induction

March 19 2024



PR55 α level is much lower in human normal tissue specimens of older individuals compared to younger individuals and inversely correlates with p16 levels. Human normal tissue specimens derived from various organs/sites were analyzed for differences in PR55 α and p16 expression by IHC. (A) Representative images of adjacent tissue sections stained with anti-PR55 α and anti-p16 antibodies. OSTSE–tonsil; GMST–seminiferous tubules; ET–thyroid; GIE–esophagus. Young, ≤ 43 y/o; Old, ≥ 68 y/o; (B) Box plot shows the H-Score quantification of PR55 α and p16 expression from adjacent sections. Credit: *Aging* (2024). DOI: 10.18632/aging.205619

A new research paper was [published](#) in *Aging* titled, "PR55 α -controlled protein phosphatase 2A inhibits p16 expression and blocks cellular senescence induction by γ -irradiation."

Cellular senescence is a permanent cell cycle arrest that both internal and external genotoxic stressors, such as telomere dysfunction and DNA damage can trigger. The execution of senescence is mainly by two pathways, p16/RB and p53/p21, which lead to CDK4/6 inhibition and RB activation to block cell cycle progression. While the regulation of p53/p21 signaling in response to DNA damage and other insults is well-defined, the regulation of the p16/RB pathway in response to various stressors remains poorly understood.

In this new study, researchers Chitra Palanivel, Lepakshe S. V. Madduri, Ashley L. Hein, Christopher B. Jenkins, Brendan T. Graff, Alison L. Camero, Sumin Zhou, Charles A. Enke, Michel M. Ouellette, and Ying Yan from the University of Nebraska Medical Center report a novel function of PR55 α , a regulatory subunit of PP2A Ser/Thr phosphatase, as a [potent inhibitor](#) of p16 expression and senescence induction by ionizing radiation (IR), such as γ -rays.

"During natural aging, there is a gradual accumulation of p16-expressing senescent cells in tissues. To investigate the significance of PR55 α in this up-regulation of p16, we compared levels of the p16 and PR55 α proteins in a panel of normal tissue specimens derived from young (≤ 43 y/o) and old (≥ 68 y/o) donors," write the researchers.

The results show that ectopic PR55 α expression in normal pancreatic cells inhibits p16 transcription, increases RB phosphorylation, and blocks IR-induced senescence. Conversely, PR55 α -knockdown by shRNA in pancreatic cancer cells elevates p16 transcription, reduces RB

phosphorylation, and triggers senescence induction after IR.

Furthermore, this PR55 α function in the regulation of p16 and senescence is p53-independent because it was unaffected by the mutational status of p53. Moreover, PR55 α only affects p16 expression but not p14 (ARF) expression, which is also transcribed from the same CDKN2A locus but from an alternative promoter. In normal human tissues, levels of p16 and PR55 α proteins were inversely correlated and mutually exclusive.

"Collectively, these results describe a novel function of PR55 α /PP2A in blocking p16/RB signaling and IR-induced [cellular senescence](#)," the authors conclude.

More information: Chitra Palanivel et al, PR55 α -controlled protein phosphatase 2A inhibits p16 expression and blocks cellular senescence induction by γ -irradiation, *Aging* (2024). [DOI: 10.18632/aging.205619](https://doi.org/10.18632/aging.205619)

Provided by Impact Journals LLC

Citation: Study: PR55 α -controlled PP2A inhibits p16 expression and blocks cellular senescence induction (2024, March 19) retrieved 29 April 2024 from <https://medicalxpress.com/news/2024-03-pr55-pp2a-inhibits-p16-blocks.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|