

Precautions needed when folks taking Ozempic, Wegovy undergo anesthesia

March 13 2024, by Carole Tanzer Miller



Nurses who specialize in anesthesia have issued [new guidelines](#) to reduce the risk that patients taking weight-loss drugs like Ozempic or Wegovy throw up during surgery.

"These medications have exploded in popularity," said Micah Walden, of the American Association of Nurse Anesthesiology (AANA) Practice Committee. "This means additional preparation for patients, anesthesia providers and the surgical team to help minimize risks of complications during a procedure."

Because [general anesthesia](#) can cause nausea, patients are usually asked to fast before surgery. But a study published last week in the journal [JAMA Surgery](#) found these weight-loss drugs — called [GLP-1 receptor agonists](#) — slow digestion, so it takes longer for food to leave the [stomach](#). That increases a patient's risk of vomiting or aspiration while under anesthesia.

GLP-1 agonists like [Ozempic](#), [Wegovy](#) (semaglutide), [Saxenda](#) (liraglutide) or [Zepbound](#) (tirzepatide) were originally designed to manage type 2 diabetes, but they have become a popular way to lose weight. They make patients feel full sooner, so they eat less.

As a precaution, the AANA committee said providers may need to do extra screenings such as an ultrasound of the patient's stomach before surgery. If the examination indicates that the stomach is not empty or the imaging is inconclusive, the surgical team may consider delaying an elective surgery or proceeding as "full stomach" to reduce the risk of vomiting and aspiration while the patient is intubated for anesthesia care, the committee said.

As they developed their recommendations, members of the practice

committee said they considered the length of time various GLP-1 medications continue to affect patients.

They recommend withholding medication for a week before surgery if patients take a weekly dose. Those on a daily dose should not use the [medication](#) on the day of procedure, the recommendations advise.

The recommendations were published March 12.

"Open communication between [patients](#) and the [surgical team](#) is important to recommendations for withholding GLP-1 agonist medications prior to [surgery](#)," Walden said in an AANA news release. "As providers, we take that information into account to perform an individualized, case-by-case assessment and create a care plan that will keep the patient safe and comfortable before, during and after the procedure."

More information: The Obesity Medicine Association has more about [popular weight-loss medications](#).

Copyright © 2024 [HealthDay](#). All rights reserved.

Citation: Precautions needed when folks taking Ozempic, Wegovy undergo anesthesia (2024, March 13) retrieved 29 April 2024 from <https://medicalxpress.com/news/2024-03-precautions-folks-ozempic-wegovy-anesthesia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.