

Replacing sugar with sweeteners can improve weight loss control over the long term in overweight adults

March 21 2024



Credit: Pixabay/CC0 Public Domain



New research being presented at this year's European Congress on Obesity (ECO) in Venice, Italy (12-15 May), suggests that replacing sugar-sweetened food and drinks with low/no energy sweetened products can help weight control for at least one year after rapid weight loss in adults, without increasing the risk of type 2 diabetes or cardiovascular disease.

The findings of a year-long randomized controlled trial involving adults with overweight and obesity and children in the overweight range from Northern, Central and Southern Europe also found that consumption of sweeteners and sweetness enhancers (S&SE) was associated with greater diet satisfaction, more positive mood, and less cravings and explicit liking for sweet foods in adults.

The results come in the wake of a 2023 systematic review of the evidence by WHO which suggested that replacing free sugars with non-sugar sweeteners did not appear to help with weight control in the long term, and could raise the risk of various health problems including type 2 diabetes and cardiovascular diseases, although the report highlighted uncertainty around the evidence between sweeteners and disease outcomes.

"Weight maintenance after weight loss is difficult to achieve, and our findings support the use of S&SEs found in many foods and beverages worldwide as alternatives to sugar-sweetened products to help manage weight control in adults with overweight," says trial lead author Professor Anne Raben from the University of Copenhagen in Denmark. "The findings also provide important information to address the controversies about potential adverse health effects. Until now, safety data have generally come from animal studies using sweetener doses far above the usual intake in humans."



S&SEs have very high sweetening power but contribute little or nothing to energy intake. Natural and synthesized sweeteners, such as aspartame, stevia, and saccharin, are commonly added to products by the food and drink industry to cut the sugar content, and are consumed by millions of people around the world every day, for example in soft drinks, desserts, and breakfast foods.

However, the safety of these food additives and their use in appetite and weight control has been hotly debated, and little is known about the long-term effects of S&SEs on eating behavior.

Sweeteners show promise for weight control

To provide more evidence, researchers from the SWEET (Sweeteners and sweetness enhancers: Prolonged effects on health, obesity and safety) project, conducted a one-year randomized controlled trial to investigate whether the prolonged use of S&SEs as part of a healthy lowsugar diet could improve weight loss maintenance after rapid weight loss, as well as benefit risk markers for type-2-diabetes and cardiovascular diseases compared to no S&SE consumption.

Participants also completed questionnaires (at the start of the trial and after 2, 6 and 12 months) to assess the effect of S&SEs on diet satisfaction, control overeating, explicit liking and implicit wanting for food, eating behavior, physical activity, and quality of life.

The trial recruited 341 adults with overweight or obesity (aged 18-65 years, 71% women, body mass index (BMI) \geq 25 kg/m²), and 38 children with overweight (6-12 years, 60% girls, BMI-for-age >85th percentile) from Denmark, Spain, Greece, and The Netherlands through web pages, social media, newspapers and registries.

During the first two months of the trial, adults were instructed to follow



a low-energy diet (the Cambridge Weight Plan) with the aim of achieving at least 5% weight loss. Children received dietary advice to maintain body weight.

In the following 10 months, participants were randomized to either a healthy diet with less than 10% of energy from added sugar allowing for foods and drinks with S&SEs (S&SEs group), or a <u>healthy diet</u> with less than 10% of energy from added sugar not allowing foods and drinks with S&SEs (sugar group).

At the start of the trial and after 2, 6, and 12 months, participants took part in clinical investigation days to measure change in weight, in BMIfor-age z-score in children (a measure of how many standard deviations a child's BMI is above or below the average BMI for their age and sex), anthropometry, and risk markers for T2D (e.g., glucose and hemoglobin A1c) and CVD (e.g., cholesterol). Participants also completed food frequency questionnaires and urine samples were collected to measure biomarkers of S&SEs, fructose and sucrose to measure compliance.

As planned, intake of sugar-rich foods and beverages declined in both groups over the trial, but significantly more in the S&SE group than in the sugar group. At the same time, intake of S&SE products increased in the S&SE group and declined in the sugar group. These results were confirmed by urinary biomarker analyses.

The intention-to-treat analyses (including 277 adults who successfully lost 5% or more body weight after the 2-month weight loss period) found that those consuming S&SEs had slightly better weight loss maintenance after 1 year than the sugar group (average weight loss of -7.2kg vs. -5.6 kg, making a difference of 1.6 kg).

Additionally, in 203 participants who completed all clinical investigations, risk markers for type 2 diabetes and cardiovascular



disease did not differ between the groups.

No differences in BMI-for-age z-score or other outcomes were noted in the 22 children who completed the trial. A larger study would be required in more children to determine any potential effect, researchers say.

Sweeteners reduce cravings for sweet foods and improve well-being in adults

In further analyses of eating behavior, researchers found that adults in the S&SE group reported greater diet satisfaction when eating out, increased positive mood, and lower craving for sweet food at 6 months than the sugar group. In contrast, adults in the sugar group had greater liking for sweet over savory foods at 6 months and 12 months. However, there were no differences between the groups in reported physical activity or quality of life.

In children, replacing food and drinks with added sugar with sweetened products had no impact on eating behavior or physical activity. However, inclusion of S&SEs appeared to benefit children with high levels of uncontrolled eating (eating in response to food palatability [tastiness] and the likelihood of over-consuming).

As co-lead author Ms. Clarissa Dakin from the Appetite Control and Energy Balance Research Group at the University of Leeds, UK, explains, "Our findings suggest that the inclusion of low/no energy sweetened products may benefit children who show high levels of uncontrolled eating. For these children, being allowed to consume S&SEs led to lower levels of uncontrolled eating at 12 months in comparison to the children who were not allowed to consume S&SE's. This benefit was not found for children with lower levels of uncontrolled



eating. Together, these findings provide important insights for the ongoing re-evaluation of food additive sweeteners by the European Food Safety Authority and other health agencies worldwide."

Professor Jason Halford, Head of the School of Psychology at the University of Leeds, UK, one of the SWEET co-authors and President of the European Association for the Study of Obesity (EASO), says, "The use of low calorie sweeteners in weight management has been questioned, in part because of the link between their use and apparent weight gain in observational studies, however, increasingly it is becoming apparent that is not the case in long term studies."

The authors note that a key strength of the study is that it is a10-month (long-term) investigation which adds to the evidence of previous shortmidterm trials by using a <u>weight loss</u> maintenance design, including both foods and drinks with S&SEs, and by using daily amounts of intervention products that are feasible and realistic in an every-day life. In addition, the trial was conducted as a multicenter trial in four countries across Europe.

Nevertheless, they note some limitations of the study, including that the drop-out rate was larger than estimated. However, with 203 completers, the statistical power for the primary outcome was 86%, which is deemed satisfactory. Additionally, they say that the children's results should be interpreted with caution because of the small number of participants. Finally, they considered S&SEs collectively without distinguishing between individual types, although this was done intentionally to reflect the variation of products in the market. Selection bias may also have been a factor, as more participants were women, had higher educational levels, and exhibited health-conscious behaviors.

This research represents the core randomized trial of the SWEET project. Prof. Jo Harrold (University of Liverpool, UK), one of the



SWEET project coordinators, explains, "The ... project has been developing and reviewing evidence on the long-term benefits and potential risks of replacing added sugar in the diet with S&SEs from multiple perspectives. The results of this long term trial support the use of S&SEs instead of sugar in food and beverages to aid <u>weight control</u>. The evidence suggests there are no grounds for concern in regard to the safety of these products on sweet preference, appetite, or glucose control.

"These results add to the body of evidence which indicates that S&SEs can play a positive role in weight management when used judiciously. Other aspects of the project have alluded to barriers that impede the development of healthier, low-energy products and a resulting need to review policy and regulatory approaches. The SWEET symposium will offer an opportunity to make industry representatives, consumers and policymakers aware of the potential benefits of S&SEs for public health and safety, obesity and sustainability."

More information: ECO 2024 Abstract:0226: Effect of sweeteners and sweetness enhancers on weight loss maintenance in adults and children with overweight or obesity – results from the SWEET multicentre, randomised, controlled, trial

ECO 2024 Abstract:0184: The impact of sweeteners on eating behaviours during a weight maintenance RCT in adults and children with overweight: SWEET project

Provided by European Association for the Study of Obesity

Citation: Replacing sugar with sweeteners can improve weight loss control over the long term in overweight adults (2024, March 21) retrieved 11 May 2024 from



https://medicalxpress.com/news/2024-03-sugar-sweeteners-weight-loss-term.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.