

Testing environmental water to monitor COVID-19 spread in unsheltered encampments

April 3 2024



Testing waterways, such as this one, located near unsheltered encampments could help monitor COVID-19 infections in a community that is often underrepresented in current surveillance methods. Credit: Edwin Oh

To better understand COVID-19's spread during the pandemic, public health officials have expanded wastewater surveillance. These efforts track SARS-CoV-2 levels and health risks among most people, but they miss people who live without shelter, a population particularly vulnerable to severe infection.

To fill this information gap, researchers reporting in [Environmental Science & Technology Letters](#) tested flood-control waterways near unsheltered encampments, finding similar transmission patterns as in the broader community and identifying previously unseen viral mutations.

In recent years, testing untreated wastewater for SARS-CoV-2 incidence and dominant viral variants, as well as other pathogens, has been vital to helping [public health officials](#) determine infectious disease transmission in [local communities](#). Yet, this monitoring only captures information on viruses shed from human feces and urine in buildings that are connected to local sewage infrastructure.

Beyond the pandemic's impact on human health, it also exacerbated socioeconomic difficulties and increased the number of people experiencing homelessness and living in open-air encampments without access to indoor bathrooms. To understand the prevalence of COVID-19 among people who live unsheltered, Edwin Oh and colleagues tested for SARS-CoV-2 in waterways near encampments outside Las Vegas from December 2021 through July 2022.

Using quantitative polymerase chain reaction, the researchers identified SARS-CoV-2 RNA in more than 25% of the samples tested from two flood-control channels. The highest detection frequency over the study period aligned with Las Vegas' first wave of omicron [variant](#) infections, as confirmed through parallel testing at a local wastewater treatment plant.

The researchers say these results suggest a similar level of transmission was occurring within the unsheltered community as it was among the general population.

Then the researchers conducted whole genome sequencing to identify the SARS-CoV-2 variants in the waterways. These samples largely contained the same variants identified in the broader community.

Deeper computational analysis of the viral sequences identified three novel viral spike protein mutations in some [waterway](#) samples, but the researchers have not yet examined what impact these mutations might have on viral function or clinical outcomes.

Regardless, the ability to detect and identify SARS-CoV-2 in environmental water samples could help improve public health measures for a community that is often underrepresented in current surveillance methods. The researchers also say monitoring waterways could warn health officials of unexpected variants circulating in the community.

More information: Environmental Surveillance of Flood Control Infrastructure Impacted by Unsheltered Individuals Leads to the Detection of SARS-CoV-2 and Novel Mutations in the Spike Gene, *Environmental Science & Technology Letters* (2024). [DOI: 10.1021/acs.estlett.3c00938](#).
pubs.acs.org/doi/abs/10.1021/acs.estlett.3c00938

Provided by American Chemical Society

Citation: Testing environmental water to monitor COVID-19 spread in unsheltered encampments

(2024, April 3) retrieved 2 May 2024 from

<https://medicalxpress.com/news/2024-04-environmental-covid-unsheltered-encampments.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.