


# High blood pressure linked to increased risk of malaria in mice

April 5 2024

## Graphical Abstract



Credit: *Function* (2024). DOI: 10.1093/function/zqae009

Hypertensive mice with abnormal red blood cells are at greater risk for developing malaria, according to a new study [published](#) ahead of print in the journal *Function*.

Until now, the link between [high blood pressure](#) and [malaria](#) has not been extensively studied. Physiologists compared hypertensive and nonhypertensive mice exposed to malaria. The findings indicate hypertensive mice displayed smaller and fewer red blood cells (microcytic anemia) that were highly resistant to hemoglobin loss.

In addition, they had increased membrane rigidity and higher levels of saturated fatty acid, indicating a higher risk of malaria. Compared to the nonhypertensive mice, the animals with elevated blood pressure experienced "significant body weight loss" and sustained parasitic infection in the blood, indicating they could not overcome malaria infection.

"Collectively, these data demonstrate that aberrant red blood cell physiology observed in hypertensive mice contributes to an increased susceptibility to malaria-associated pathology," the researchers wrote.

**More information:** Mrunmayee R Kandalgaoonkar et al, Hypertension Increases Susceptibility to Experimental Malaria in Mice, *Function* (2024). [DOI: 10.1093/function/zqae009](https://doi.org/10.1093/function/zqae009)

Provided by American Physiological Society

Citation: High blood pressure linked to increased risk of malaria in mice (2024, April 5) retrieved 2 May 2024 from

<https://medicalxpress.com/news/2024-04-high-blood-pressure-linked-malaria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.