

International experts agree on standards for catheter ablation of atrial fibrillation

April 8 2024



Credit: Pixabay/CC0 Public Domain

An international consensus statement on how to treat atrial fibrillation with catheter or surgical ablation has been [published](#) in *EP Europace* and presented at [EHRA 2024](#).

Lead author Dr. Stylianos Tzeis said, "Ablation is the most effective way to prevent recurrences of atrial fibrillation and delay progression to more advanced forms. Pioneering techniques have emerged since the previous [consensus in 2017](#), requiring new advice on who should receive this procedure and how to perform it in the safest and most effective manner."

Atrial fibrillation is the most common cardiac arrhythmia, affecting 2% of individuals worldwide. The number of people with the disorder has risen three-fold over the last 50 years and is expected to increase further—mainly due to population aging and unhealthy lifestyles—reaching 14 million by 2060 in the European Union alone. It is estimated that approximately one in three White and one in five Black individuals will develop atrial fibrillation during their lifetime.

Atrial fibrillation has a significant impact on people's lives. Symptoms include palpitations, fatigue, shortness of breath, dizziness, and fainting. Compared to their peers without the condition, people with atrial fibrillation are twice as likely to be admitted to hospital, five times more likely to have a stroke, three times more likely to develop [heart failure](#), and twice as likely to die prematurely.

Catheter ablation of atrial fibrillation is an important treatment option for maintaining a normal heart rhythm and reducing arrhythmia-related symptoms. Ablation is a procedure which uses burning, freezing, or high energy electrical pulses (called electroporation) to inactivate the electrical triggers causing abnormal rhythms. It is typically a one- to two-hour, minimally invasive procedure in which a catheter is inserted into a blood vessel in the groin and advanced into the heart, with patients going home the same or next day.

The recent document sets out best practice standards and practical advice on which patients should receive ablation, how to perform the

procedure, and how to manage patients before, during and afterwards. A [healthy lifestyle](#) is advised to reduce the risk of developing atrial fibrillation and to lower the number of recurrences.

Catheter ablation is beneficial as first-line treatment in patients with symptoms and recurrent atrial fibrillation. It is also beneficial in symptomatic patients with recurrent paroxysmal or persistent atrial fibrillation who are resistant or intolerant to at least one antiarrhythmic drug. Patients with atrial fibrillation and heart failure, a condition in which the heart muscle cannot pump enough blood to meet the body's needs, benefit from ablation to alleviate symptoms, increase the heart's strength, and improve their prognosis.

Before the procedure, patients should have an electrocardiogram (ECG) and echocardiogram (ultrasound of the heart) to check the heart's rhythm and function. Some patients need computed tomography or transesophageal echocardiography to exclude the presence of a blood clot in the heart. Blood thinners should not be interrupted before [catheter ablation](#) and patients should receive specific instructions from their health care provider on continuing their medication.

After the procedure, all patients should receive anticoagulation for at least two months. Prolonged anticoagulation may be required depending on the risk of blood clots. Antiarrhythmic drugs are advised for some patients to prevent arrhythmia recurrences early after the procedure. Complications following catheter ablation are uncommon but can include stroke, pericardial tamponade (fluid accumulation around the heart), [transient ischemic attack](#), and vascular complications at the access site in the groin. The risk of death is extremely low (0.05–0.1%).

Dr. Tzeis said, "Technological innovations have made catheter ablation safer and more effective than ever before. For example, we now have pulsed field ablation which uses high energy [electrical pulses](#) to treat

[atrial fibrillation](#) without damaging the nerves or esophagus. We can also use intracardiac echocardiography to guide and facilitate the procedure with real-time visualization of the cardiac structures being treated."

More information: Alireza Sepehri Shamloo et al, European Heart Rhythm Association (EHRA)/Heart Rhythm Society (HRS)/Asia Pacific Heart Rhythm Society (APHRS)/Latin American Heart Rhythm Society (LAHRS) Expert Consensus Statement on Catheter and Surgical Ablation of Atrial Fibrillation, *EP Europace* (2024). [DOI: 10.1093/europace/euae043](#)

Provided by European Society of Cardiology

Citation: International experts agree on standards for catheter ablation of atrial fibrillation (2024, April 8) retrieved 17 May 2024 from <https://medicalxpress.com/news/2024-04-international-experts-standards-catheter-ablation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--