

Majority of people with heart disease consume too much sodium, research finds

April 2 2024



Credit: Pixabay/CC0 Public Domain

Individuals with heart disease stand to gain the most from a low-sodium diet but, on average, consume over twice the recommended daily sodium intake, according to a study being presented at the [American College of Cardiology's Annual Scientific Session](#).

Sodium is an essential nutrient, but consuming too much can raise blood pressure, which damages blood vessels and forces the heart to work harder. Excess sodium can also cause the body to retain fluid, exacerbating conditions like heart failure.

The current U.S. Dietary Guidelines put out by the U.S. Department of Agriculture recommends most adults limit their sodium intake to less than 2,300 mg/day, which is equivalent to about 1 teaspoon of table salt. For individuals with cardiovascular diseases, the limit is even lower at 1,500 mg/day, according to [guideline recommendations](#) from the ACC and the American Heart Association.

This new study found that among a sample of more than 3,100 people with [heart disease](#), 89% consumed more than the recommended daily maximum of 1,500 mg of sodium and, on average, study participants consumed more than twice this amount. Limiting sodium intake is a fundamental lifestyle modification shown to reduce the likelihood of subsequent major adverse cardiovascular events, researchers said.

Their findings underscore the challenges many people face in keeping within recommended sodium limits, regardless of other factors such as socioeconomic status.

"Estimating sodium quantities in a meal can be challenging," said Elsie Kodjoe, MD, MPH, an internal medicine resident at Piedmont Athens Regional Hospital in Athens, Georgia and the study's lead author. "Food

labels aid in dietary sodium estimation by providing sodium quantities in packaged food. Yet, adhering to a low sodium diet remains challenging even for individuals with cardiovascular disease who have a strong incentive to adhere."

The study used data from patients diagnosed with a heart attack, stroke, [heart failure](#), [coronary artery disease](#) or angina who participated in the National Health and Nutrition Examination Survey (NHANES) between 2009–2018.

Researchers estimated sodium intake based on questionnaires in which participants were asked to report everything they had consumed in 24 hours. According to the results, study participants with cardiovascular disease consumed an average of 3,096 mg of sodium per day, which is slightly lower than the national average of 3,400 mg/day reported by the U.S. Centers for Disease Control and Prevention.

"The relatively small difference in sodium intake suggests that people with cardiovascular disease are not limiting their intake very much compared with the general population and are also consuming more than double what is recommended," Kodjoe said. "To make it easier for patients to adhere to [dietary guidelines](#), we need to find more practical ways for the general public to estimate dietary sodium levels or perhaps consider a reduction in the sodium content of the food we consume right from the source."

The researchers also compared sodium intake among people in different socioeconomic groups, but they did not find any significant differences between wealthier and less affluent participants after accounting for age, sex, race and educational attainment.

Individuals can take proactive measures to lower their sodium intake, Kodjoe said. This includes preparing more meals at home where they

have greater control over the sodium content and paying close attention to [food labels](#), particularly targeting foods with sodium levels of 140 mg or less per serving. Researchers suggested that better education around the benefits of limiting sodium could also help motivate more people to follow the recommendations.

"Cardiovascular disease is real, and it is the number one cause of morbidity and mortality worldwide according to the World Health Organization," Kodjoe said. "Adhering to sodium guidelines is one of the easier strategies individuals could readily adopt to reduce hospitalizations, [health care costs](#), morbidity and mortality associated with cardiovascular disease."

One limitation of the study is that sodium intake was estimated based on food recall questionnaires, rather than 24-hour urine sodium measurements, which is considered the gold standard method. NHANES has included 24-hour urine sodium measurements in its data gathering methods in recent survey cycles, so future studies using this data could provide a more accurate assessment of [sodium intake](#) among people with cardiovascular disease.

More information: Kodjoe will present the study, "The Impact of Income to Poverty Ratio on Sodium Intake Among Adults with Cardiovascular Disease," on Sunday, April 7, 2024.

For more information on reducing sodium, visit [CardioSmart.org/Sodium](https://www.heart.org/sod-smart)

Provided by American College of Cardiology

Citation: Majority of people with heart disease consume too much sodium, research finds (2024,

April 2) retrieved 2 May 2024 from <https://medicalxpress.com/news/2024-04-majority-people-heart-disease-consume.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.