

Optimizing differentiation protocols and experimental assays to study patient-specific astrocytes

April 16 2024



Credit: Kyoto University

In a joint effort with RIKEN BioResource Research Center and Mitsubishi Tanabe Pharma Corporation, a team of researchers led by Dr. Haruhisa Inoue (Professor, Department of Cell Growth and Differentiation, CiRA) improved upon a differentiation protocol to generate astrocytes from patient-derived induced pluripotent stem (iPS) cells for modeling an astrocyte-associated disease.

Their study is <u>published</u> in the *Journal of Cellular and Molecular Medicine*.



Astrocytes, one of two major types of supporting cells in the central nervous system (i.e., brain and <u>spinal cord</u>), are increasingly recognized for their involvement in many human diseases including stroke and Alzheimer's disease. As such, it is critical to determine how they are affected in the pathophysiological state and whether they could be targeted for treatment and cure.

The research team had previously established a "feeder-free" protocol to generate <u>astrocytes</u> from iPS cells (iPSCs), thus enabling disease modeling and drug discovery. However, establishing high-throughput screening platforms based on human iPSC-derived astrocytes requires robust and reproducible functional assays with the potential for large-scale operations.

Since such experimental pipelines to analyze various astrocytic functions are currently unavailable, therapeutic development using iPSC-derived astrocytes has been limited. Therefore, in this study, the researchers primarily focused on validating several functional assays designed to assess vital biological processes that astrocytes are involved in and could dramatically impact disease progression: cytokine release, calcium flux, autophagy, and <u>cell migration</u>.

Neuroinflammation, particularly the release of inflammatory cytokines from astrocytes, can be detrimental to brain cells during <u>disease</u> <u>progression</u> or following injuries. Therefore, the researchers determined whether iPSC-derived astrocytes are responsive to inflammatory stimuli such as tumor necrosis factor-alpha (TNF- α) and interleukin 1-beta (IL-1 β). Following TNF- α or IL-1 β treatment, the research team detected cytokines released into the media from iPSC-derived astrocytes, thus demonstrating their responsiveness to inflammatory stimulation.

Astrocytic calcium homeostasis is especially crucial to neuronal



functions and survival. The researchers examined whether iPSC-derived astrocytes can undergo ATP-induced calcium uptake and detected dose-dependent increases in calcium influx of the astrocytes.

Autophagy, one of the principal degradative pathways in cells, is enhanced by nutrient starvation and often compromised in neurodegenerative diseases, not only in neurons but also in astrocytes. The researchers measured autophagy using a fluorescent probe in normal and starvation conditions and observed an increase in autophagy in iPSCderived astrocytes after starvation, reduced when it was chemically blocked.

Astrocytes migrate to lesion sites to mediate wound healing and tissue repair in response to traumatic injuries and other stimuli. Because these processes can drastically affect functional recovery, the research team also assessed whether iPSC-derived astrocytes properly migrate in response to stimulation. Through the scratch assay, in which cells are scratched off the surface of a cell culture plate, they observed both iPSCderived astrocytes and primary astrocytes to similarly migrate into the injured region.

Lastly, having confirmed that iPSC-derived astrocytes behave similarly to primary astrocytes, the research team next generated astrocytes from two iPSC lines, one established from a healthy volunteer and another from an Alexander disease (AxD) patient carrying a mutation in the GFAP gene. The researchers observed a significant increase in GFAPpositive aggregates in AxD iPSC-derived astrocytes.

In summary, the research team validated iPSC-derived astrocytes to function similarly to primary astrocytes. Crucially, their work reveals the potential of using iPSC-derived astrocytes as an in vitro model of disease or injury. Furthermore, the assays examined by the researchers are ripe for adaptation to establish high-throughput screening pipelines for



identifying chemical or <u>genetic factors</u> affecting astrocytic functions under patient-specific disease conditions.

Altogether, the series of experiments in this study paves the way forward for drug discovery against various astrocyte-associated pathologies using iPSC-derived astrocytes.

More information: Hideki Nonaka et al, Induced pluripotent stem cell-based assays recapture multiple properties of human astrocytes, *Journal of Cellular and Molecular Medicine* (2024). DOI: 10.1111/jcmm.18214

Provided by Kyoto University

Citation: Optimizing differentiation protocols and experimental assays to study patient-specific astrocytes (2024, April 16) retrieved 17 May 2024 from <u>https://medicalxpress.com/news/2024-04-optimizing-differentiation-protocols-experimental-assays.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.