



Proximal femur bionic nail offers new hope for elderly patients with intertrochanteric femur fractures

April 1 2024



Schematic of (a) the proximal femur bionic nail (PFBN) and (b) its modified

version for preventing bone nonunion, Yingze nail. Credit: Kaixuan Zhang, Wei Chen, Yingze Zhang

In a new study [published](#) in *Engineering*, academician Yingze Zhang's research team introduces the Proximal Femur Bionic Nail (PFBN) as a potential panacea for unstable intertrochanteric femur fractures in the elderly.

With an [aging population](#), intertrochanteric [femur](#) fractures have become a critical public health concern, prompting a surge in research focused on trauma orthopedics.

Traditional internal fixation techniques have often fallen short in addressing the mechanical challenges posed by these fractures, leading to prolonged recovery times and increased complication rates for patients. However, the development of the PFBN, incorporating Zhang's N triangle theory to provide triangular supporting fixation, represents a [paradigm shift](#) in the treatment of these fractures.

The key innovation of the PFBN lies in its ability to significantly reduce the occurrence of complications, leading to improved patient outcomes and faster recovery times. Through the use of an equivalent biomechanical model, the research team demonstrated that the PFBN outperforms existing solutions—such as the proximal femoral nail antirotation (PFNA) and InterTan—by dramatically decreasing maximum strain in the proximal femur.

Furthermore, the PFBN has been shown to regulate the local mechanical environment in the proximal femur, creating a state of physiological

overload that favors post-operative recovery for elderly patients with intertrochanteric femur fractures. This promising outcome offers hope for a transformative approach to internal fixation, with the potential to revolutionize treatment not only for this specific type of fracture but for other fractures as well, based on elastic deformation.

Zhang's research team concludes that the PFBN holds the potential to revolutionize the clinical treatment of intertrochanteric femur fractures in the elderly, enabling patients to avoid prolonged bed rest and exercise, and reducing the likelihood of complications. The implications of this study extend beyond the specific application of the PFBN, pointing toward a promising avenue for further advancements in the development of next-generation internal fixation systems.

The findings of this research have far-reaching implications, offering a glimmer of hope for the growing population of elderly individuals at risk of intertrochanteric femur fractures. As the [global community](#) grapples with the challenges posed by an aging population, the emergence of the PFBN represents a significant advancement with the potential to enhance the quality of life for countless individuals worldwide.

More information: Kaixuan Zhang et al, Proximal Femur Bionic Nail (PFBN): A Panacea for Unstable Intertrochanteric Femur Fracture, *Engineering* (2024). [DOI: 10.1016/j.eng.2024.01.010](https://doi.org/10.1016/j.eng.2024.01.010)

Provided by Engineering

Citation: Proximal femur bionic nail offers new hope for elderly patients with intertrochanteric femur fractures (2024, April 1) retrieved 21 May 2024 from <https://medicalxpress.com/news/2024-04-proximal-femur-bionic-elderly-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.