

## Study provides first stratification of the risk of developing dilated cardiomyopathy in symptom-free genetic carriers

April 22 2024



Cardiac MRI of a patient with dilated cardiomyopathy (left) and an individual without heart disease (right). In the patient with dilated cardiomyopathy, the size of the heart is increased and its pumping capacity is reduced. Credit: CNIC



Dilated cardiomyopathy is the most frequent cause of heart failure in young people and is the leading cause of heart transplantation. In this disease, the heart enlarges and reduces its capacity to pump blood. People with dilated cardiomyopathy are at high risk for arrhythmias and sudden death.

In approximately 30%–40% of people with dilated cardiomyopathy, the disease is caused by a genetic mutation. When a genetic cause is identified, the patient's <u>family members</u> can be studied to determine if they have also inherited the altered gene.

Family members who are carriers of the genetic mutation are at risk for developing the disease in the future, and it is recommended that they undergo regular check-ups so that any signs of the disease can be detected as soon as they appear and pharmacological treatments initiated.

Nevertheless, it is not known if all family members carrying the culprit mutation will go on to develop the disease, at what age this is more likely to happen, and if there are factors that could help to identify those individuals who are likely to show signs of disease over the short term and who would therefore benefit from closer monitoring.

The new study, led by Dr. Pablo García-Pavía, a researcher at the Centro Nacional de Investigaciones Cardiovasculares (CNIC), a group leader in the Spanish cardiovascular research network (CIBERCV), and a <u>cardiologist</u> at Hospital Puerta de Hierro, provides the first stratification of the risk of developing dilated cardiomyopathy among symptom-free genetic carriers of the disease.

A total of 25 Spanish hospitals participated in the study, which is published in the *Journal of the American College of Cardiology*. Data were collected from more than 779 genetic carriers, from 300 families, who had shown no signs of the disease before entering the study.



The researchers found that, after a median follow-up of 37 months, almost 11% of the genetic carriers had developed the disease.

The study also showed that the appearance of the disease was dependent on the specific type of genetic mutation present.

Moreover, said Dr. García-Pavía, "We found that patients who develop the disease tended to share certain characteristics, such as older age, electrocardiogram alterations, and an enlarged heart with a weaker pumping action, albeit within the normal range."

Dr. García-Pavía added that "the risk for developing the disease was also higher in patients with mutations in sarcomere genes or in those whose hearts showed signs of fibrosis detected by magnetic resonance imaging."

First author of the study, Dr. Eva Cabrera, a cardiologist at Hospital Puerta de Hierro, said, "These results will allow us to offer more personalized care to genetic carriers with no disease symptoms."

Currently, all genetic carriers receive a similar care strategy, consisting of annual or twice-yearly check-ups. "Now, we can schedule check-ups to meet the specific needs of the individual patient: more spaced out for patients without <u>risk factors</u>, and more frequent for those at higher risk," said Dr. Cabrera.

**More information:** Penetrance of Dilated Cardiomyopathy in genotype positive relatives, *Journal of the American College of Cardiology* (2024).

Provided by Centro Nacional de Investigaciones Cardiovasculares Carlos



## III (F.S.P.)

Citation: Study provides first stratification of the risk of developing dilated cardiomyopathy in symptom-free genetic carriers (2024, April 22) retrieved 27 June 2024 from <u>https://medicalxpress.com/news/2024-04-stratification-dilated-cardiomyopathy-symptom-free.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.