

Study shows aerobic exercise performed in the evening benefits elderly hypertensives more than morning exercise

May 17 2024, by Maria Fernanda Ziegler



Credit: Wellness Gallery Catalyst Foundation from Pexels



Aerobic training is known to regulate blood pressure more effectively when practiced in the evening than in the morning. Researchers who conducted a study of elderly patients at the University of São Paulo's School of Physical Education and Sports (EEFE-USP) in Brazil concluded that evening exercise is better for blood pressure regulation thanks to improved cardiovascular control by the autonomic nervous system via a mechanism known as baroreflex sensitivity. The study is <u>published</u> in *The Journal of Physiology*.

"There are multiple mechanisms to regulate blood pressure, and although morning training was beneficial, only evening training improved shortterm control of blood pressure by enhancing baroreflex sensitivity. This is important because baroreflex control has a positive effect on <u>blood</u> <u>pressure regulation</u>, and there aren't any medications to modulate the mechanism," Leandro Campos de Brito, first author of the article, told.

The study was part of Brito's postdoctoral research project, which is supported by FAPESP and supervised by Cláudia Lúcia de Moraes Forjaz, a professor at EEFE-USP.

In the study, 23 <u>elderly patients</u> diagnosed and treated for hypertension were randomly allocated into two groups: morning training and evening training. Both groups trained for ten weeks on a stationary bicycle at moderate intensity, with three 45-minute sessions per week.

Key cardiovascular parameters were analyzed, such as systolic and diastolic blood pressure, and <u>heart rate</u> after ten minutes' rest. The data was collected before and at least three days after the volunteers completed the ten weeks of training.

The researchers also monitored mechanisms pertaining to the autonomic



nervous system (which controls breathing, heart rate, blood pressure, digestion, and other involuntary bodily functions), such as muscle sympathetic nerve activity (which regulates peripheral blood flow via contraction and relaxation of blood vessels in <u>muscle tissue</u>) and sympathetic baroreflex sensitivity (assessing control of blood pressure via alterations to muscle sympathetic nerve activity).

In the evening training group, all four parameters analyzed were found to improve: systolic and <u>diastolic blood pressure</u>, sympathetic baroreflex sensitivity, and muscle sympathetic nerve activity. In the morning training group, no improvements were detected in muscle sympathetic nerve activity, <u>systolic blood pressure</u> or sympathetic baroreflex sensitivity.

"Evening training was more effective in terms of improving cardiovascular autonomic regulation and lowering blood pressure. This can be partly explained as due to an improvement in baroreflex sensitivity and a reduction of muscle <u>sympathetic nerve activity</u>, which increased in the evening.

"For now, all we know is that baroreflex control is the decisive factor, from the cardiovascular standpoint at least, to make evening training more beneficial than morning training, since it induces the other benefits analyzed. However, much remains to be done in this regard in order to obtain a better understanding of the mechanisms involved," said Brito, who is currently a professor at Oregon Health & Science University's Oregon Institute of Occupational Health Sciences in the United States, and continues to investigate the topic via circadian rhythm studies.

Baroreflex sensitivity regulates each heartbeat interval and controls autonomic activity throughout the organism. "It's a mechanism that involves sensitive fibers and deformations in the walls of arteries in specific places, such as the aortic arch and carotid body.



"When blood pressure falls, this region warns the brain region that controls the autonomic nervous system, which in turn signals the heart to beat faster and tells the arteries to contract more strongly. If blood pressure rises, it warns the heart to beat more slowly and tells the arteries to contract less. In other words, it modulates arterial pressure beat by beat," Brito explained.

In previous studies, the EEFE-USP research group showed that evening aerobic training reduced blood pressure more effectively than morning training in hypertensive men, and that the more effective response to evening training in terms of blood pressure control was accompanied by a greater reduction in systemic vascular resistance and systolic pressure variability.

"Replication of the results obtained in previous studies and in different groups of hypertensive patients, associated with the use of more precise techniques to evaluate the main outcomes, has strengthened our conclusion that aerobic exercise performed in the evening is more beneficial to the autonomic nervous system in patients with hypertension. This can be especially important for those with resistance to treatment with medication," Brito said.

More information: Leandro C. Brito et al, Evening but not morning aerobic training improves sympathetic activity and baroreflex sensitivity in elderly patients with treated hypertension, *The Journal of Physiology* (2024). DOI: 10.1113/JP285966

Provided by FAPESP

Citation: Study shows aerobic exercise performed in the evening benefits elderly hypertensives more than morning exercise (2024, May 17) retrieved 21 June 2024 from



https://medicalxpress.com/news/2024-05-aerobic-evening-benefits-elderly-hypertensives.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.