

Study finds association between neighborhood deprivation and DNA methylation in an autopsy cohort

May 9 2024



Manhattan (A) and QQ plot (B) from the EWAS of DNAm with the ADI.



Adjusted for race, sex, educational attainment, age at death, APOE genotype, cell type, and post-mortem interval. Bonferroni-threshold: $0.05/789889 = 6.33e^{-8}$ (λ =0.94). Credit: *Aging* (2024). DOI: 10.18632/aging.205764

A new research paper was <u>published</u> in *Aging*, titled, "The association between neighborhood deprivation and DNA methylation in an autopsy cohort."

Previous research has found that living in a disadvantaged neighborhood is associated with poor health outcomes. Living in disadvantaged neighborhoods may alter inflammation and immune response in the body, which could be reflected in <u>epigenetic mechanisms</u> such as DNA methylation (DNAm).

In this new study, researchers from Emory University, University of British Columbia, BC Children's Hospital Research Institute, Centre for Molecular Medicine and Therapeutics, and Atlanta VA Medical Center used robust linear regression models to conduct an epigenome-wide association study examining the association between neighborhood deprivation (Area Deprivation Index; ADI), and DNAm in brain tissue from 159 donors enrolled in the Emory Goizueta Alzheimer's Disease Research Center (Georgia, U.S.).

"We found one CpG site (cg26514961, gene PLXNC1) significantly associated with ADI after controlling for covariates and multiple testing (p-value = $5.0e^{-8}$)," write the researchers.

Effect modification by APOE ε 4 was statistically significant for the top ten CpG sites from the EWAS of ADI, indicating that the observed associations between ADI and DNAm were mainly driven by donors who carried at least one APOE ε 4 allele. Four of the top ten CpG sites



showed a significant concordance between brain tissue and tissues that are easily accessible in living individuals (blood, buccal cells, saliva), including DNAm in cg26514961 (PLXNC1).

This study identified one CpG site (cg26514961, PLXNC1 gene) that was significantly associated with neighborhood deprivation in <u>brain</u> <u>tissue</u>. PLXNC1 is related to <u>immune response</u>, which may be one biological pathway to how neighborhood conditions affect health.

"The concordance between brain and other tissues for our top CpG sites could make them potential candidates for biomarkers in living individuals," the authors conclude.

More information: Lindsay Pett et al, The association between neighborhood deprivation and DNA methylation in an autopsy cohort, *Aging* (2024). DOI: 10.18632/aging.205764

Provided by Impact Journals LLC

Citation: Study finds association between neighborhood deprivation and DNA methylation in an autopsy cohort (2024, May 9) retrieved 28 June 2024 from <u>https://medicalxpress.com/news/2024-05-association-neighborhood-deprivation-dna-methylation.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.