

Chronic stress during adolescence may reduce fertility in adulthood

May 10 2024



Credit: Unsplash/CC0 Public Domain



Male rats exposed to moderate and repeated stress during adolescence may have reduced fertility, according to research presented at the <u>26th</u> <u>European Congress of Endocrinology</u>, held 11-14 May, in Stockholm. This study sheds light on the harmful effects that early-life stress has on health and could help to uncover future prevention strategies for children and adolescents.

While <u>hormone levels</u> fluctuate pathologically—especially during life stages such as puberty—stress can cause too much or too little of a hormone in the bloodstream. This hormonal imbalance negatively impacts puberty and the reproduction system, with effects on libido, ovulation function and sperm cell production. However, the long-term reproductive effects of chronic stress in adolescents are largely unknown.

In this study, researchers from the V.P. Komisarenko Institute of Endocrinology and Metabolism of the National Academy of Medical Science of Ukraine, examined male and <u>female rats</u>, aged 6 months, after individually placing about half of them in enclosed spaces for one hour every morning for two weeks during adolescence (30–45 days old).

They compared the rats who had been exposed to these stressful conditions to the control group and found that chronic stress during puberty delayed <u>sexual maturity</u> in females, and males gained weight more slowly. In adult males, sperm count fell by 25.9%, some sperm were abnormally shaped and sluggish or became immobile, and the breathing process by which sperm cells derive energy slowed down. Additionally, males had almost two times lower levels of corticosterone—the main stress hormone in rats, equivalent to cortisol in humans.



"Our work is the first to report to show that even moderate and repetitive stress in adolescence has a long-lasting negative impact on the endocrine system of reproduction and adaptation of the body to changing living conditions," said lead investigator, Professor Aleksander Reznikov.

"Our results make it possible to predict the development of anomalies in reproduction and bodily adaptation systems and are the basis for finding methods for their prevention."

Professor Reznikov added, "We discovered for the first time that <u>lipid</u> <u>peroxidation</u> (a process in which oxidants like free radicals attack lipid membranes of cells and eventually damage them) in the ovaries and testes was significantly increased. This, however, needs further investigation."

Provided by European Society of Endocrinology

Citation: Chronic stress during adolescence may reduce fertility in adulthood (2024, May 10) retrieved 16 June 2024 from <u>https://medicalxpress.com/news/2024-05-chronic-stress-adolescence-fertility-adulthood.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.