


Study develops light-activated drug targeting receptors in the brain that induce sleep

May 23 2024



Activation of NAc A_{2A}R by focal injection of adenosine or stimulation of astrocytes induces SWS. Credit: *Nature Communications* (2024). DOI: 10.1038/s41467-024-47964-4

Adenosine A_{2A} receptors (A_{2A}R) in the nucleus accumbens of the brain play an important role in regulating sleep and motivation, but until now, no drugs have been able to selectively modulate their function without affecting other organs or brain areas. In a recent study, a light-activated allosteric modulator of A_{2A}R was developed and successfully used to remotely induce sleep by selective light irradiation of the nucleus accumbens in mice.

The nucleus accumbens plays a pivotal role in motivational behavior and sleep regulation, modulated by adenosine A_{2A} receptors (A_{2A}R). Hence, selective A_{2A}R regulation within this brain region could control sleep and motivation. However, A_{2A}Rs are distributed across various organs, including the heart, posing challenges for precise brain-specific modulation without genetic interventions.

A research team led by Professor Michael Lazarus and Associate Professor Tsuyoshi Saitoh (TRiSTAR Fellow) from the Institute of Medicine and the International Institute for Integrative Sleep Medicine (WPI-IIIS) at the University of Tsukuba delved into optochemistry. [The study](#) is published in *Nature Communications*.

They aimed to develop a novel light-sensitive drug that enhances extracellular adenosine activity. By administering this drug to [mice](#) and selectively irradiating the [nucleus accumbens](#) with light, they succeeded in inducing sleep artificially without [genetic modification](#) for the first

time.

Conventional photosensitive drugs have faced hurdles in mammals and other living organisms due to problems such as phototoxicity caused by [ultraviolet light](#), blood-brain barrier permeability, and photoreaction efficiency.

The newly developed photosensitive drug overcomes these issues, showcasing optochemistry's potential in developing drugs targeting A_{2A}R in the brain and regulating brain function by targeting other central drug receptors.

More information: Koustav Roy et al, Optochemical control of slow-wave sleep in the nucleus accumbens of male mice by a photoactivatable allosteric modulator of adenosine A_{2A} receptors, *Nature Communications* (2024). [DOI: 10.1038/s41467-024-47964-4](https://doi.org/10.1038/s41467-024-47964-4)

Provided by University of Tsukuba

Citation: Study develops light-activated drug targeting receptors in the brain that induce sleep (2024, May 23) retrieved 16 June 2024 from <https://medicalxpress.com/news/2024-05-drug-receptors-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.