

Study shows extending interval between colonoscopies feasible after negative result

May 3 2024, by Elana Gotkine



For individuals without a family history of colorectal cancer (CRC), increasing the interval between colonoscopies for those with a first colonoscopy with negative findings seems safe and can avoid unnecessary colonoscopies, according to a study [published](#) online May 2 in *JAMA Oncology*.

Qunfeng Liang, from the German Cancer Research Center in Heidelberg, and colleagues assessed how many years after a first colonoscopy with findings negative for CRC a second colonoscopy can be performed.

The exposed group included individuals without a family history of CRC who had a first colonoscopy with findings negative for CRC at age 45 to 69 years between 1990 and 2016, while the [control group](#) included matched individuals who did not have a colonoscopy during follow-up or who underwent colonoscopy resulting in CRC diagnosis (110,074 and 1,981,332 individuals, respectively).

The researchers found that 484 incident CRCs and 112 CRC-specific deaths occurred during up to 29 years of follow-up of individuals with a first colonoscopy with findings negative for CRC.

For 15 years, the [risks](#) for CRC and CRC-specific [death](#) were significantly lower in the exposed group than in matched controls. The 10-year standardized incidence ratio was 0.72 at 15 years after a first colonoscopy with negative findings, and the 10-year standardized mortality ratio was 0.55.

Extending the colonoscopy screening interval from 10 to 15 years could miss the early detection of two CRC cases and [prevention](#) of one CRC-specific death per 1,000 individuals and could potentially avert 1,000 colonoscopies.

"A longer interval between [colonoscopy](#) screenings could be beneficial in avoiding unnecessary invasive examinations," the authors write.

More information: Qunfeng Liang et al, Longer Interval Between First Colonoscopy With Negative Findings for Colorectal Cancer and Repeat Colonoscopy, *JAMA Oncology* (2024). [DOI](#):

[10.1001/jamaoncol.2024.0827](https://doi.org/10.1001/jamaoncol.2024.0827)

Rashid N. Lui et al, The Interval for Screening Colonoscopy—Is 15 the New 10?, *JAMA Oncology* (2024). DOI: [10.1001/jamaoncol.2024.0249](https://doi.org/10.1001/jamaoncol.2024.0249)

© 2024 [HealthDay](https://www.healthday.com/). All rights reserved.

Citation: Study shows extending interval between colonoscopies feasible after negative result (2024, May 3) retrieved 30 June 2024 from <https://medicalxpress.com/news/2024-05-interval-colonosopies-feasible-negative-result.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.