

Study investigates whether Weo electrolyzed water can modulate phenotype of senescent cells

May 22 2024



WEW decreases cellular senescence in human fibroblasts and cancer cells. Credit: *Aging* (2024). DOI: 10.18632/aging.205789



A new research paper titled "Cell type-dependent modulation of senescence features using Weo electrolyzed water" has been <u>published</u> in *Aging*.

Electrolyzed-reduced water has powerful antioxidant properties with constituents that scavenge <u>reactive oxygen species</u> (ROS), which are known to be produced by several intrinsic and extrinsic processes. When there is an imbalance between ROS production and antioxidant defenses, <u>oxidative stress</u> occurs. Persistent oxidative stress leads to <u>cellular</u> <u>senescence</u>, an important hallmark of aging, and is involved in several age-related conditions and illnesses.

In this new study, researchers Brenda L. Court-Vazquez, Shirley A. Arroyo-Vizcarrondo, Jonathan A. Poli, Lara Nyman, Kelly Halderman, Anthony Ginter, and Pierre-Yves Desprez from Weo LLC and California Pacific Medical Center investigated whether Weo electrolyzed water (WEW) could modulate the phenotype of senescent cells.

"The focus of this study was to utilize two different cell types, human normal fibroblasts and human breast <u>cancer cells</u>, to investigate the impact of Weo electrolyzed water (WEW) on markers of cellular senescence, inflammation, and stress response genes," the researchers explain.

The researchers compared normal human lung fibroblasts (BJ) and breast cancer cells (T47D) treated with <u>hydrogen peroxide</u> (H_2O_2) to induce senescence. They assessed the molecular impact of WEW on markers of cellular senescence, senescence-associated secretory



phenotype (SASP) factors, and stress response genes. Treatment with WEW modulated markers of cellular senescence, such as the senescenceassociated β -galactosidase (SA- β -gal) activity, EdU incorporation and p21 expression, similarly in both cell types.

However, WEW modulated the expression of SASP factors and stress response genes in a cell type-dependent and opposite fashion, significantly decreasing them in BJ cells, while stimulating their expression in T47D cells. Reduction in the expression of SASP factors and stress-related genes in BJ cells suggests that WEW acts as a protective factor, thereby reducing oxidative stress in normal cells, while making cancer cells more sensitive to the effects of cellular stress, thus increasing their elimination and consequently reducing their deleterious effects.

"In conclusion, we have shown here that the new technology developed by Weo, WEW, could attenuate the overall process of cellular senescence in both normal BJ fibroblasts and cancer T47D cells," the researchers state.

More information: Brenda L. Court-Vazquez et al, Cell typedependent modulation of senescence features using Weo electrolyzed water, *Aging* (2024). <u>DOI: 10.18632/aging.205789</u>

Provided by Impact Journals LLC

Citation: Study investigates whether Weo electrolyzed water can modulate phenotype of senescent cells (2024, May 22) retrieved 22 June 2024 from https://medicalxpress.com/news/2024-05-weo-electrolyzed-modulate-phenotype-senescent.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.