

## Alarming study unveils how forever chemicals transfer from mothers to newborns

June 24 2024



Box-plots of concentrations of PFAS with > 50% detection in maternal serum across trimesters (T1–T3), cord serum or breast milk (ng/mL). Credit: *Eco-Environment & Health* (2024). DOI: 10.1016/j.eehl.2024.04.007

Polyfluoroalkyl substances (PFAS) are a class of chemicals extensively used in consumer goods production due to their hydrophobic and oleophobic properties and stability. However, their persistence in the environment and bioaccumulation in living organisms have sparked



concerns about potential health effects. Previous studies have linked PFAS exposure to various adverse outcomes, including developmental issues in children.

A <u>recent study</u> published in *Eco-Environment & Health* on May 8, 2024, has highlighted the levels, health risks, and <u>transport protein</u> binding capabilities of per- and <u>polyfluoroalkyl substances</u> (PFAS) in <u>early life</u> based on the Shanghai Maternal-Child Pairs Cohort.

Found in maternal serum, cord serum, and breast milk, these synthetic chemicals pose potential health risks for infants. Led by a team from School of Public Health at Fudan University, the researchers meticulously analyzed the transfer mechanisms and impacts of these persistent chemicals, providing crucial insights into their pervasive presence from pregnancy to lactation.

Employing high-performance liquid chromatography-tandem mass spectrometry, the study analyzed 16 types of PFAS in 1,076 motherchild pairs. It revealed the detection rates and median concentrations of perfluorooctane sulfonate (PFOS), <u>perfluorooctanoic acid</u> (PFOA), and 6:2 Cl-PFESA, with PFOS most prevalent in maternal serum.

Notably, placental transfer efficiency of PFAS was higher than breastfeeding transfer, suggesting that these chemicals are more likely to cross the placenta and accumulate in the fetus. Additionally, the research utilized molecular docking to simulate the binding of PFAS to transport proteins, potentially influencing their distribution and transport within the body.

These findings underscore the need for tighter PFAS regulations and further research into their environmental and health effects.

Ms. Yaqi Xu, lead author of the study, states, "Our findings are crucial



for developing strategies to protect infants from the potentially harmful effects of PFAS exposure. Understanding the pathways and risks associated with these chemicals can lead to better regulatory policies and protective measures for the most susceptible among us."

The implications of this research are profound, particularly for public health policies and infant safety. By identifying specific PFAS compounds more likely to transfer through the placenta and into <u>breast</u> <u>milk</u>, preventive measures can be more effectively targeted. Moreover, the study's findings could influence future guidelines on the use of PFAS-containing products by <u>pregnant women</u> and nursing mothers.

**More information:** Yaqi Xu et al, Early-life exposure to per- and polyfluoroalkyl substances: Analysis of levels, health risk and binding abilities to transport proteins, *Eco-Environment & Health* (2024). DOI: 10.1016/j.eehl.2024.04.007

Provided by Chinese Academy of Sciences

Citation: Alarming study unveils how forever chemicals transfer from mothers to newborns (2024, June 24) retrieved 12 September 2024 from <u>https://medicalxpress.com/news/2024-06-alarming-unveils-chemicals-mothers-newborns.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.