

Over-the-counter supplement found to improve walking for peripheral artery disease patients

June 18 2024, by Eric Hamilton



Credit: Yaroslav Shuraev from Pexels

The over-the-counter supplement nicotinamide riboside, a form of vitamin B3, increased the walking endurance of patients with peripheral



artery disease, a chronic leg condition for which there are few effective treatments.

In a preliminary, randomized, double-blind clinical trial led by Northwestern University and University of Florida scientists, patients who took nicotinamide riboside daily for six months increased their timed walking distance by more than 57 feet, compared to participants who took a placebo. As expected, walking speed declined in those who took a placebo, because peripheral artery disease causes progressive declines in walking performance.

"This is a signal that nicotinamide riboside could help these patients," said Christiaan Leeuwenburgh, Ph.D., a UF professor of physiology and aging and senior author of the clinical trial report. "We are hoping to conduct a larger follow-up trial to verify our findings."

Leeuwenburgh, whose research specializes in anti-aging treatments, collaborated with Mary M. McDermott, M.D., a physician and professor of medicine at Northwestern University and an expert in peripheral artery disease. With a large team of collaborators, Leeuwenburgh and McDermott <u>published their findings</u> June 13 in the journal *Nature Communications*.

The scientists recruited 90 people with an average age of 71 who had peripheral artery disease, or PAD, to test the effects of nicotinamide riboside. The supplement is increasingly popular as an anti-aging treatment—sales exceeded \$60 million in 2022 in the U.S. alone—but there has been scant evidence of any benefit in healthy people. Nicotinamide riboside is a precursor for the essential compound NAD, which plays roles in the body related to energy generation, improved blood flow and DNA repair.

Because PAD is associated with problems generating energy within



muscle cells, McDermott and Leeuwenburgh thought that nicotinamide riboside, by improving energy generation, could help improve walking in people with the disease.

And indeed that's what they found. Participants taking the supplement walked an average of 23 feet more in a six-minute walking test after six months, while those taking a placebo walked 34 feet less. Those who took at least 75% of the pills they were supposed to take performed even better, adding more than 100 feet to their walking distance, compared to people who took a placebo.

(The researchers also tested if resveratrol, a compound best known for being in red wine, could boost the effects of nicotinamide riboside; they found no additional benefits.)

PAD affects more than 8.5 million Americans over the age of 40. Caused by the buildup of fatty deposits in arteries, and associated with diabetes and smoking, the disease reduces blood flow to the limbs, especially the legs. Walking often becomes painful, and the disease typically causes declines in walking ability over time. Supervised walking exercise is first line therapy for PAD, but most people with the condition do not have access to supervised exercise.

In addition to a larger trial focused on patients suffering from PAD, Leeuwenburgh hopes to test the effects of <u>nicotinamide riboside</u> on walking performance in healthy older adults.

"We need to test it on a healthy older population before we recommend healthy people take it," he said.

More information: Mary M. McDermott et al, Nicotinamide riboside for peripheral artery disease: the NICE randomized clinical trial, *Nature Communications* (2024). DOI: 10.1038/s41467-024-49092-5



Provided by University of Florida

Citation: Over-the-counter supplement found to improve walking for peripheral artery disease patients (2024, June 18) retrieved 29 June 2024 from https://medicalxpress.com/news/2024-06-counter-supplement-peripheral-artery-disease.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.