


What is the neural mechanism behind helping someone at your own cost?

June 12 2024



Helping task structure. Top: two screen shots taken from a video showing the confederate receiving an electrical shock on the hand and manifesting its painfulness through facial reactions and a video showing the confederate receiving a slap on her right hand. Middle: trial structure. Bottom: the experiment consisted of two Face and two Hand blocks were presented of 15 trials each. Credit: *Scientific Reports* (2024). DOI: 10.1038/s41598-024-62422-3

Using a unique setup, researchers from the Social Brain Lab at the Netherlands Institute for Neuroscience have researched the neural mechanism behind a universal dilemma: deciding whether to help someone else even when it involves a personal sacrifice.

We often have opportunities to give up something we care about to help others. What brain mechanisms help us make those decisions, and why do some people help more than others?

Over the years, philosophers and scientists have suggested that the extent to which a person empathizes with the distress of others influences their willingness to help.

To explore this hypothesis experimentally, Kalliopi Ioumpa and Selene Gallo, under the supervision of Christian Keysers and Valeria Gazzola, investigated pro-social behavior—voluntary behavior intended to benefit others—in self-reported mirror-pain synaesthetes. Mirror-pain synaesthetes report feeling pain on their own body whenever they're witnessing pain on someone else. If sharing the pain of others is indeed a key motivator to help others, these individuals should be particularly generous.

"To better understand why some people help more than others, in the past, our lab had performed experiments with participants that are at the lower extreme of empathy, including individuals with psychopathy. This time, we wanted to look at the opposite extreme: people that feel the pain of others so much more, that it feels like the pain is on their own body," explains Keysers.

"In the literature, there was some notion that these participants feel such a localized pain when witnessing that of others, because their brain activated somatosensory cortices—regions normally only involved in the sense of touch on your own body. Would that influence their

donations?".

The conflict scenario

Studying a realistic conflict in a controlled lab-setting requires a creative approach in which participants feel that their decisions have real consequences both for themselves and for someone else. "If you just ask people whether they are generous or not, most of them like to say yes" Keysers says with a smile.

"To reveal real differences between people, you have to create a situation in which the distress of others feels real, and where helping has a real cost for participants," Gazzola emphasizes.

In their study, they created such a conflict by making participants believe that they were participating in the study together with another participant. In reality, the other participant was an actor, which kept each experiment consistent.

In the experiment the two "participants" were allocated two different roles: one, the actor, who was filmed while receiving painful stimulations on their hand; the other, the participant, who received real money and could donate some of this money to alleviate some of the other participant's pain.

"All participants believed that the other was really receiving the painful stimulation on their hand in ways that depended on their donations," Ioumpa explains. "In reality we pre-recorded the actor receiving these painful stimulations ahead of time, so that all participants saw the exact same stimuli."

Because participants made their decisions in the fMRI scanner, the researchers could analyze the [brain areas](#) related to their decisions, and

how they differed between those that feel pain on their own body when they see the pain of others, and those that do not. Ioumpa says, "We wanted to explore how much different participants would help and what the underlying neural mechanisms would be."

Mirror-pain synaesthetes and prosocial behavior

Compared to control participants, the self-reported mirror-pain synaesthetes donated significantly more money to alleviate the other participant's pain—giving credence to the belief that the more you feel the pain of others, the more you are willing to give up to help them.

As expected, the researchers also found significant differences between the two groups when looking at the brain mechanism "self-reported mirror pain synesthetes activated their secondary somatosensory cortex more. This brain area is involved in sensory experiences such as touch or pain," Ioumpa explains.

"So while most of us just feel generally uncomfortable when we see others in pain, mirror touch synesthetes have a higher activation in their somatosensory cortex, as if they are experiencing pain in the corresponding body parts themselves, and they are more motivated to help," adds Keysers.

Towards more prosocial behavior

After each experiment, participants are told the truth about the other participant being an actor, and they receive the amount of money they choose to keep.

Ioumpa says, "We know that someone's level of empathy can change. People can learn to enhance it but also to shut it down. Consider doctors

for example, they tend to become less empathetic over the years because they need to block their empathy."

Gazzola adds, "And as I explain in my TEDx talk, we can all modulate how empathic we want to be in a particular situation. Empathy is not a fixed trait, it's something we have control over, and that, as we show here, can influence how prosocial we decide to be."

"This study helps us understand prosocial behavior better as well, as its links to empathy. We show that a group of highly empathic individuals, that somatically feels the [pain](#) of others, also demonstrates enhanced prosocial behavior," Ioumpa explains. The researchers hope that their work can inspire interventions to enhance empathy and prosocial behavior in the future.

More information: Kalliopi Ioumpa et al, Neural mechanisms of costly helping in the general population and mirror-pain synesthetes, *Scientific Reports* (2024). [DOI: 10.1038/s41598-024-62422-3](https://doi.org/10.1038/s41598-024-62422-3)

Provided by Netherlands Institute for Neuroscience - KNAW

Citation: What is the neural mechanism behind helping someone at your own cost? (2024, June 12) retrieved 18 June 2024 from <https://medicalxpress.com/news/2024-06-neural-mechanism.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
