

## **People who exercise more may have a lower risk of ALS**

June 26 2024



Credit: Pixabay/CC0 Public Domain

Moderate levels of physical activity and fitness may be linked to a reduced risk of amyotrophic lateral sclerosis (ALS) later in life, according to a new study published in *Neurology*. However, the study



only found an association between physical activity and risk of ALS in male participants, not female participants.

ALS is a rare, progressive neurodegenerative disease that affects <u>nerve</u> <u>cells</u> in the brain and the spinal cord. People with ALS lose the ability to initiate and control muscle movement, which often leads to total paralysis and death. The average life span after diagnosis is two to five years.

"The diagnosis of prominent athletes with ALS at young ages has sparked the uncomfortable idea that higher physical activity could be tied to developing ALS," said study author Anders Myhre Vaage, MD, of Akershus University Hospital in Norway. "There have been conflicting findings on levels of physical activity, fitness and ALS risk. Our study found that for men, living a more active lifestyle could be linked to a reduced risk of ALS more than 30 years later."

For the study, researchers looked at 373,696 people in Norway with an average age of 41. They were followed for an average of 27 years.

Of the total participants, 504 people developed ALS. Of those who developed ALS, 59% were male participants.

Participants recorded their level of physical activity for the past year into one of four categories: sedentary; a minimum of four hours per week of walking or cycling; a minimum of four hours per week of recreational sports or heavy gardening; or participation in hard training or sports competitions regularly, several times a week. Due to few participants with the highest level of physical activity, researchers combined the third and fourth categories into one high activity group.

Researchers found that of the 41,898 male participants that had the highest level of physical activity, 63 developed ALS; of the 76,769 male



participants with the intermediate level of physical activity, 131 developed ALS; and of the 29,468 male participants with the lowest level of physical activity, 68 developed ALS.

After adjusting for other factors that could affect the risk of ALS, such as smoking and body mass index, researchers found that for <u>male</u> <u>participants</u>, when compared to those with the lowest level of physical activity, those with moderate levels of physical activity had a 29% lower risk of ALS and those with high levels of physical activity had a 41% lower risk of ALS.

Researchers also looked at resting heart rate. Men in the lowest of four categories of resting <u>heart rate</u>, which indicates good physical fitness, had a 32% reduced risk of ALS compared to those with higher resting heart rates.

"Our findings show that for men, not only do moderate to high levels of physical activity and fitness not increase the risk of ALS, but that it may be protective against the disease," Myhre Vaage said. "Future studies of the connection between ALS and exercise are needed to consider sex differences and higher or professional athlete physical activity levels."

A limitation of the study was that the <u>physical activity</u> questionnaire was completed only at one specific time during the study, so it may not have captured the participants' exercise levels over the nearly 30-year span of the study.

Provided by American Academy of Neurology

Citation: People who exercise more may have a lower risk of ALS (2024, June 26) retrieved 30 June 2024 from <u>https://medicalxpress.com/news/2024-06-people-als.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.