


Finding epilepsy hotspots before surgery: A faster, non-invasive approach

August 27 2024, by Nancy Fliesler



Credit: Pixabay/CC0 Public Domain

Neurosurgery for patients with drug-resistant epilepsy requires locating the precise brain areas that are generating the seizures. Typically, patients undergo seven to 10 days of invasive intracranial EEG

monitoring, with electrodes surgically implanted inside the brain through one or more skull openings to capture seizure activity as it happens.

Eleonora Tamilia, Ph.D., directs the Epilepsy Monitoring Unit Signal and Data Science Program within the Epilepsy Center at Boston Children's Hospital. Her team has piloted a much briefer method for mapping seizure zones.

Not only is it noninvasive, but can it provide information a traditional EEG reading cannot. It combines standard scalp EEG readings with MRI data on [brain structure](#) to identify connections between different areas of the brain and uses machine learning to locate the most epileptogenic brain regions.

"Using [computational tools](#), we can reconstruct cortical activity that the eye cannot catch and understand how different regions are functionally connected," explains Tamilia, who is also part of the Fetal-Neonatal Developmental Science Center (FNNDSC). "If a seizure starts in one [region](#) of the cortex, it's likely to spread to another network it connects to. Even regions that are far apart may fire together."

Predicting seizure zones

As described in [Epilepsia](#), Tamilia and her colleagues retrospectively analyzed about five minutes of scalp EEG data from 50 patients with drug-resistant epilepsy who had neurosurgery.

Adding MRI data and using machine learning algorithms, they were able to define functional cortical networks—even during times when no epileptiform activity was visible to the naked eye and even when the MRI showed no obvious brain abnormalities.

"When there were spikes on the EEG, our predictions were more

precise, but we could still get an estimate without them," Tamilia says.

During epileptiform activity, the algorithm achieved 75% accuracy (91% sensitivity, 74% specificity) in identifying the surgically targeted seizure zones. During quiet periods, it achieved 62% accuracy.

The algorithm was less likely to match the targeted zones in patients who continued to have seizures after surgery, suggesting that the implemented surgical plan did not actually pinpoint the seizure focus. "This suggests that the plan could have been changed," Tamilia says. "In cases of unsuccessful surgery, the model indicated epileptogenic regions that were not completely removed."

The tool could also indicate that the epilepsy area is too broad to resect, and that patients may instead benefit from other palliative treatments such as neuromodulation, she adds.

The researchers now hope to validate their findings in a larger, prospective cohort and define which [patients](#) with drug-resistant epilepsy are most likely to benefit from surgery. Because their technique is brief and noninvasive, it could be used early in the course of disease, Tamilia says.

This could enable some children to have epilepsy surgery sooner rather than later, avoiding some of [epilepsy](#)'s neurodevelopmental outcomes.

More information: Georgios Ntolkeras et al, Interictal EEG source connectivity to localize the epileptogenic zone in patients with drug-resistant epilepsy: A machine learning approach, *Epilepsia* (2024). [DOI: 10.1111/epi.17898](https://doi.org/10.1111/epi.17898)

Provided by Children's Hospital Boston

Citation: Finding epilepsy hotspots before surgery: A faster, non-invasive approach (2024, August 27) retrieved 27 August 2024 from <https://medicalxpress.com/news/2024-08-epilepsy-hotspots-surgery-faster-invasive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.