


Inflammation may link COVID-19 severity to the onset of neurological symptoms

August 26 2024



Representative paired neuroimaging and chest CT findings from COVID-19 patients presenting neurological symptoms. Credit: *Brain, Behavior, & Immunity - Health* (2024). DOI: 10.1016/j.bbih.2024.100805

COVID-19 is primarily known for its effects on the respiratory system, but its consequences go far beyond that. A recent study, [published](#) in the journal *Brain, Behavior, & Immunity—Health* and conducted at the D'Or Institute for Research and Education (IDOR), has revealed molecular changes that may underlie the neurological symptoms exhibited by some patients affected by the disease, highlighting the importance of better understanding these still poorly known potential consequences of COVID-19.

COVID-19 continues to be a concerning disease even after the end of its pandemic phase. In the first half of 2024 alone, it was responsible for over 3,000 deaths in Brazil. Additionally, scientific literature has been extensively documenting the deleterious effects of the infection even after patients have recovered, a condition now known as "long COVID."

Long COVID refers to a range of persistent symptoms that remain or emerge after the acute phase of the disease. Even after recovering from respiratory symptoms, patients may continue to face significant challenges, especially concerning neurological health. A significant portion of COVID-19 survivors, even those who had mild cases, may experience cognitive decline and difficulty concentrating for extended periods after the infection. This makes it important to study how the disease affects the brain even in the acute phase, as this may provide clues about these neurological sequelae.

During the infection, the most common [neurological symptoms](#) are

headaches, fatigue, loss of smell, and even more severe complications such as strokes and encephalitis. Investigating these manifestations is crucial, as we still know little about the mechanisms leading to these complications and how they develop.

How the study was conducted

Seeking biomarkers that could provide clues about neuroinflammatory processes in COVID-19, IDOR researchers analyzed data from patients confirmed to have COVID-19 who were hospitalized in the Rede D'Or network between April and November 2020.

The sample included 35 patients aged between 26 and 87 years, divided between moderate and severe cases, all of whom presented significant neurological symptoms during the acute COVID-19 infection. The data were collected from [medical records](#) and included imaging exams (MRI and CT scans), blood tests, and [cerebrospinal fluid](#) (CSF) analysis—a fluid that surrounds the brain and spinal cord. Ten CSF samples from uninfected patients served as a control group.

Findings reveal significant brain inflammation

The analysis revealed that most patients had at least one comorbidity, with 65.7% having two or more. About 85.7% of patients exhibited neurological symptoms at the time of hospital admission, a clinical picture that was even more pronounced than respiratory symptoms.

Imaging exams showed that 28.6% of patients had focal or diffuse brain changes associated with COVID-19, including demyelinating lesions, encephalitis, and stroke.

Blood tests indicated that 66% of patients showed signs of an

exacerbated inflammatory response. Proteomic analyses of the CSF revealed an altered protein pattern compared to the controls, with 116 significantly dysregulated proteins related to the immune system and metabolic processes.

Pro-inflammatory cytokines are associated with disease severity

The levels of two [pro-inflammatory cytokines](#), IL-6 and TNF α , were elevated in the CSF of COVID-19 patients, with IL-6 being particularly higher in severe cases. These cytokines are associated with the severity of the disease and the changes observed in imaging exams.

Dr. Fernanda Aragão, a postdoctoral researcher at IDOR and the study's first author, comments that the research is one of the first to connect imaging exams and neurological symptoms with neuroinflammatory biomarkers capable of reflecting the severity of acute disease, a complication that remains difficult to predict.

Despite the severity distinction found from these biomarkers, the researcher emphasizes that neuroinflammation is independent of [disease severity](#) and may be one of the main causes of neurological disorders associated with COVID-19. She points out that even patients with milder cases showed significant changes in the CSF, suggesting that the body's inflammatory response may affect the brain in ways not yet fully understood.

"This study reveals that neuroinflammation is a common factor in neurological cases of the disease, even in patients with diverse conditions, whether moderate or severe. Identifying these inflammatory markers that connect COVID-19 severity and neuroimaging changes could be very important for developing therapies aimed at both treating

acute COVID-19 infections and addressing the persistent effects of what is known as long COVID," adds the author.

Implications for long-term treatment and monitoring

These findings highlight the need for long-term monitoring of patients who have had COVID-19, especially those at risk of developing persistent neurological complications. Better understanding these mechanisms can help develop more effective treatment and prevention strategies in the future.

IDOR's research provides valuable insights into the neurological impacts of COVID-19 and paves the way for future studies to explore these findings more deeply and in larger populations.

Thus, continued investigation into the neurological consequences of COVID-19 is presented as a crucial investment, especially with the evolution of new variants and the implementation of vaccination programs, to ensure that the long-term effects of the pandemic are adequately understood and addressed.

More information: Fernanda G.Q. Barros-Aragão et al, Changes in neuroinflammatory biomarkers correlate with disease severity and neuroimaging alterations in patients with COVID-19 neurological complications, *Brain, Behavior, & Immunity - Health* (2024). [DOI: 10.1016/j.bbih.2024.100805](https://doi.org/10.1016/j.bbih.2024.100805)

Provided by D'Or Institute for Research and Education

Citation: Inflammation may link COVID-19 severity to the onset of neurological symptoms (2024, August 26) retrieved 27 August 2024 from

<https://medicalxpress.com/news/2024-08-inflammation-link-covid-severity-onset.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.