


Lung nodules seen in a high percentage of non-smokers

August 13 2024



Representative chest low-dose CT images show lung nodules (arrows) and were included in the study. (A) A solid lung nodule, 82 mm^3 , in a 77-year-old male participant. (B) A clinically relevant nodule, 174 mm^3 , in a 55-year-old female participant. (C) An actionable nodule, 422 mm^3 , in a 79-year-old female

participant. (D) A typical perifissural nodule, 183 mm³, in a 46-year-old male participant. Credit: Radiological Society of North America

A new study of more than 10,000 non-smoking adults found that solid lung nodules were present in a considerable portion of study participants. Non-smokers are traditionally thought to be at low risk for lung nodules and lung cancer. The results of the study were published in *Radiology*.

Incidental [lung nodules](#) are common findings on chest CT and in [high-risk groups](#) are more likely to be a sign of early-stage lung cancer. Because most previous research on the prevalence and size of lung [nodules](#) has typically been derived from lung cancer screening studies in heavy smokers, nodule management recommendations are mostly based on this patient population.

Thus, current nodule follow-up guidelines which are mostly based on high-risk patient populations may lead to many unnecessary follow-up examinations in low-risk individuals with incidental pulmonary nodules.

"This study is groundbreaking as it provides the first comprehensive analysis of the prevalence and size distribution of solid lung nodules in a population-based Northern European non-smoking [cohort](#)," said senior author Rozemarijn Vliegenthart, M.D., Ph.D., radiologist and professor of cardiothoracic imaging at the University Medical Center Groningen and the University of Groningen in the Netherlands.

"Unlike prior studies that predominantly targeted high-risk lung cancer screening cohorts or Asian cohorts, this research yields fundamental data for the general non-smoking population in northern Europe."

The study included 10,431 never- or former smokers aged 45 and over

from the Imaging in LifeLines (ImaLife) study, the first population-based imaging study designed to establish the reference values of imaging biomarkers for early stages of coronary artery disease, lung cancer and [chronic obstructive pulmonary disease](#) in a general population of mostly non-smokers.

Of the participants in the current study, 56.6% were women (median age 60.4 years), 46.1% were never smokers and 53.9% were former smokers.

"By including a large cohort of non-smoking men and women aged 45 years and above, this research offers insights into the prevalence and characteristics of lung nodules in a population group that has not been studied before," Dr. Vliegenthart said.

Participants underwent a low-dose CT exam of the chest. Seven trained readers registered the presence and size of solid lung nodules. Along with the prevalence and size of the lung nodules, they noted clinically relevant and actionable nodules.

At least one lung nodule was found in 4,377 (42.0%) of the participants (47.5% of men; 37.7% of women). The prevalence of lung nodules increased with age. Clinically-relevant lung nodules (nodules measuring 6 to 8 mm) were present in 11.1% of participants, with the prevalence again increasing with age. Men were more likely than women to have lung nodules, and men were more likely than women to have multiple nodules.

"Our study revealed the presence of clinically relevant nodules in 11.1% of a non-smoking cohort, traditionally considered low risk," Dr. Vliegenthart said. "This was higher than we expected and even similar to the prevalence reported in high-risk populations of smokers."

In recent years, the proportion of smoking individuals in Western populations has been declining due to the success of smoking cessation programs, Dr. Vliegenthart noted.

"This shift makes our study, which provides foundational and comprehensive data on lung nodules in non-smokers, even more critical," she said.

The results imply that, given the increasing numbers of chest CT scans for various clinical indications, these incidental nodules will increase the number of follow-up scans and workups when using current nodule follow-up guidelines. As the patients are followed in the LifeLines cohort over time, we can gather more data to determine if lower risk patient population may need altered follow-up management.

"We know that the incidence of lung cancer in this population (LifeLines cohort), is very low (0.3%), suggesting that most of the clinically relevant and even actionable nodules in a non-smoking cohort are benign," Dr. Vliegenthart said.

"Future data on [lung cancer](#) diagnosis in ImaLife participants with clinically relevant and actionable nodules may help to optimize nodule management recommendations for individuals considered at low risk."

More information: Distribution of Solid Lung Nodules Presence and Size by Age and Sex in a Northern European Nonsmoking Population, *Radiology* (2024). doi.org/10.1148/radiol.231436

Provided by Radiological Society of North America

Citation: Lung nodules seen in a high percentage of non-smokers (2024, August 13) retrieved 14

August 2024 from <https://medicalxpress.com/news/2024-08-lung-nodules-high-percentage-smokers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.