

Ovarian cancer is hard to detect—focusing on these four symptoms can help with diagnosis

August 18 2024, by Jenny Doust



Credit: Unsplash/CC0 Public Domain

Ovarian cancers are often found when they are already advanced and hard to treat.

Researchers [have long believed](#) this was because women first experienced [symptoms](#) when ovarian cancer was already well-established. Symptoms can also be [hard to identify](#) as they're vague and similar to other conditions.

But [a new study](#) shows promising signs ovarian cancer can be detected in its early stages. The study targeted women with four specific symptoms—bloating, abdominal pain, needing to pee frequently, and feeling full quickly—and put them on a [fast track](#) to see a specialist.

As a result, even the most aggressive forms of ovarian cancer could be detected in their early stages.

So what did the study find? And what could it mean for detecting—and treating—ovarian cancer more quickly?

Why is ovarian cancer hard to detect early?

Ovarian cancer [cannot be detected](#) via [cervical cancer screening](#) (which used to be called a [pap smear](#)) and pelvic exams [aren't useful](#) as a screening test.

Current [Australian guidelines](#) recommend women get tested for ovarian cancer if they have symptoms for [more than a month](#). But many of the [symptoms](#)—such as tiredness, constipation and changes in menstruation—are vague and overlap with other common illnesses.

This makes early detection a challenge. But it is crucial—a woman's [chances of surviving ovarian cancer](#) are associated with how advanced the cancer is when she is diagnosed.

If the cancer is still confined to the original site with no spread, the five-year survival rate is 92%. But over half of women diagnosed with

ovarian cancer first present when the cancer has [already metastatised](#), meaning it has spread to other parts of the body.

If the cancer has spread to nearby lymph nodes, the survival rate is reduced to 72%. If the cancer has already metastasized and spread to distant sites at the time of diagnosis, the rate is only 31%.

There are mixed findings on whether detecting ovarian cancer earlier leads to better survival rates. For example, a trial in the UK that screened more than 200,000 women [failed to reduce deaths](#).

That study screened the general public, rather than relying on self-reported symptoms. The new study suggests asking women to look for specific symptoms can lead to earlier diagnosis, meaning treatment can start more quickly.

What did the new study look at?

Between June 2015 and July 2022, the researchers recruited 2,596 women aged between 16 and 90 from 24 hospitals across the UK.

They were asked to monitor for these four symptoms:

- persistent abdominal distension (women often refer to this as bloating)
- feeling full shortly after starting to eat and/or loss of appetite
- pelvic or [abdominal pain](#) (which can feel like indigestion)
- needing to urinate urgently or more often.

Women who reported at least one of four symptoms persistently or frequently were put on a [fast-track pathway](#). That means they were sent to see a gynecologist within two weeks. The fast track pathway has been used in the UK since 2011, but is not specifically part of Australia's

guidelines.

Some 1,741 participants were put on this fast track. First, they did a [blood test](#) that measured the cancer antigen 125 (CA125). If a woman's CA125 level was abnormal, she was sent for an internal vaginal ultrasound.

What did they find?

The study indicates this process is better at detecting ovarian cancer than general screening of people who don't have symptoms. Some 12% of women on the fast-track pathway were diagnosed with some kind of ovarian cancer.

A total of 6.8% of fast-tracked patients were diagnosed with high-grade serous ovarian cancer. It is the most aggressive form of cancer and responsible for 90% of ovarian cancer deaths.

Out of those women with the most aggressive form, one in four were diagnosed when the cancer was still in its early stages. That is important because it allowed treatment of the most lethal cancer before it had spread significantly through the body.

There were some promising signs in treating those with this aggressive form. The majority (95%) had surgery and three quarters (77%) had chemotherapy. Complete cytoreduction—meaning all of the cancer appears to have been removed—was achieved in six women out of ten (61%).

It's a promising sign that there may be ways to "catch" and target ovarian cancer before it is well-established in the body.

What does this mean for detection?

The study's findings suggest this method of early testing and referral for the symptoms leads to earlier detection of ovarian cancer. This may also improve outcomes, although the study did not track survival rates.

It also points to the importance of public awareness about symptoms.

Clinicians should be able to recognize all of the ways ovarian cancer can present, including vague symptoms like general fatigue.

But empowering members of the general public to recognize a narrower set of four symptoms can help trigger testing, detection and treatment of ovarian cancer earlier than we thought.

This could also save GPs advising every woman who has general tiredness or constipation to undergo an ovarian cancer test, making testing and treatment more targeted and efficient.

Many women remain [unaware of the symptoms](#) of ovarian cancer. This study shows recognizing them may help early detection and treatment.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Ovarian cancer is hard to detect—focusing on these four symptoms can help with diagnosis (2024, August 18) retrieved 18 August 2024 from <https://medicalxpress.com/news/2024-08-ovarian-cancer-hard-focusing-symptoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.