


# Retrospective study explores mitral valve surgery outcomes in atrial functional mitral regurgitation

August 19 2024



Researchers provide new insights into the causes of Atrial functional Mitral Regurgitation, a rare heart condition. Credit: Nobuyuki Kagiyama of Juntendo University

Mitral regurgitation (MR) is a serious heart condition that often requires corrective surgery. It is characterized by the backflow or "regurgitation"

of blood from the heart's left ventricle into the left atrium.

Atrial functional [mitral regurgitation](#) (AFMR), characterized by normal mitral valve (MV) function and left ventricular function but with atrial dilation and defects in the ring-like structure that supports the MV leaflets (mitral annulus), poses diagnostic and therapeutic challenges due to its distinct pathophysiological mechanisms.

Historically, studies on AFMR have been limited by small sample sizes and single-center data, leading to variability in reported prevalence and [clinical outcomes](#).

To address this gap, a team of researchers led by Associate Professor Nobuyuki Kagiya from the Department of Cardiovascular Biology and Medicine at Juntendo University Graduate School of Medicine, Japan, collaborated with Tomohiro Kaneko and Tohru Minamino, also from the same department, alongside Minoru Tabata from the Department of Cardiovascular Surgery at Juntendo University School of Medicine, and Yukio Abe from the Department of Cardiology at Osaka City General Hospital.

Together, they conducted a [retrospective study](#) analyzing echocardiographic reports of patients who underwent transthoracic echocardiography. Their findings were [published](#) on August 15, 2024, in *JAMA Network Open*.

Called the REVEAL-AFMR study, the team analyzed a vast dataset comprising 225,163 echocardiographic reports from 177,235 patients who underwent transthoracic echocardiography in 2019. Through rigorous screening and validation processes, the study identified 1,007 cases of AFMR, accounting for 11.4% of all patients with MR in the study cohort.

The study's findings shed light on several pivotal aspects of AFMR. Firstly, AFMR was found to predominantly affect [elderly patients](#) burdened with conditions like [atrial fibrillation](#), enlarged left atria, and stiffened left ventricles.

Secondly, the study's evaluation of MV surgery outcomes upon a 3-year follow-up in 113 patients revealed a marked reduction in MR severity post-operation, correlating with substantially better clinical outcomes, including smaller rates of heart failure hospitalizations and all-cause mortality.

These findings are the first report of outcomes after MV surgery in AFMR, and will be an important basis of decision-making in management strategies of AFMR.

"The study showed that patients with AFMR generally have poor prognoses, but those who underwent mitral valve surgery had significantly better outcomes. Notably, the surgical group had a three-year event rate of 18.3% compared to 33.3% in the non-surgical group," observes Kagiya.

"These findings underscore the importance of understanding AFMR and its [treatment options](#). We believe this study provides critical insights into the real-world characteristics and treatment strategies for AFMR, paving the way for future trials to explore causal relationships and optimize patient care."

Although patients with AFMR are older, the study revealed that younger patients were most likely to receive corrective mitral valve surgery. Compared to those who remained on medical therapy, patients who underwent surgery also had more severe MR, had greater left atrium volume and suffered heart failure more frequently.

Although further research is required to prove the relationships definitively, the general trends in AFMR and outcomes of corrective mitral valve surgery in patients uncovered by this study are expected to guide clinicians in improved decision-making. The knowledge generated may also stimulate more scientific investigations into the origins of AFMR.

"This research addresses the significant challenge of underdiagnosis and undertreatment of AFMR by providing robust data on its clinical characteristics and outcomes," Dr. Kagiya explains.

"Health care providers can use this information to develop better screening protocols and treatment strategies, potentially improving the quality of life and survival rates for patients with AFMR. Additionally, this study could influence health care policies and [resource allocation](#) to ensure that patients with AFMR receive the attention and care they need."

"Overall, our research contributes to a deeper understanding of AFMR and paves the way for future studies and clinical trials aimed at optimizing management and treatment for affected patients," he concludes.

**More information:** Nobuyuki Kagiya et al, Clinical Outcomes of Mitral Valve Surgery in Atrial Functional Mitral Regurgitation in the REVEAL-AFMR Registry, *JAMA Network Open* (2024). [DOI: 10.1001/jamanetworkopen.2024.28032](https://doi.org/10.1001/jamanetworkopen.2024.28032)

Provided by Juntendo University Research Promotion Center

Citation: Retrospective study explores mitral valve surgery outcomes in atrial functional mitral

regurgitation (2024, August 19) retrieved 20 August 2024 from  
<https://medicalxpress.com/news/2024-08-retrospective-explores-mitral-valve-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.