

Doxorubicin + trabectedin tied to increased survival in metastatic leiomyosarcoma

September 6 2024, by Elana Gotkine



For patients with metastatic or surgically unresectable leiomyosarcoma, combination therapy with doxorubicin and trabectedin is associated with improved overall and progression-free survival compared with



doxorubicin alone, according to a study <u>published</u> online Sept. 4 in the *New England Journal of Medicine*.

Patricia Pautier, M.D., from the Institut Gustave-Roussy in Villejuif, France, and colleagues conducted a Phase III trial involving 150 patients with metastatic or unresectable leiomyosarcoma who had not received chemotherapy.

Patients were randomly assigned to <u>doxorubicin</u> (six cycles) or doxorubicin plus trabectedin (six cycles), with continued trabectedin as maintenance therapy in patients in the doxorubicin-trabectedin group without disease progression. After six cycles of therapy, surgery was allowed to resect residual disease in each group.

The researchers found that at a median follow-up of 55 months, 47 and 60 patients in the doxorubicin-trabectedin and doxorubicin groups, respectively, had died. Median overall survival was longer in the doxorubicin-trabectedin group versus the doxorubicin group (33 versus 24 months; adjusted hazard ratio for death, 0.65).

Progression-free survival was longer in the doxorubicin-trabectedin group versus the doxorubicin group, in a finding consistent with earlier reports (12 versus six months; adjusted hazard ratio for progression or death, 0.37).

"The trial results support the use of doxorubicin plus trabectedin for the first-line treatment of advanced or metastatic leiomyosarcomas, offering hope for improved outcomes in this challenging disease area," the authors write.

More information: Patricia Pautier et al, Doxorubicin–Trabectedin with Trabectedin Maintenance in Leiomyosarcoma, *New England Journal of Medicine* (2024). DOI: 10.1056/NEJMoa2403394



Robert S. Benjamin, A Late-Arriving but Welcome Advance in Sarcoma Therapy, *New England Journal of Medicine* (2024). DOI: <u>10.1056/NEJMe2407116</u>

Copyright © 2024 <u>HealthDay</u>. All rights reserved.

Citation: Doxorubicin + trabectedin tied to increased survival in metastatic leiomyosarcoma (2024, September 6) retrieved 11 September 2024 from https://medicalxpress.com/news/2024-09-doxorubicin-trabectedin-survival-metastaticleiomyosarcoma.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.