

# Optical coherence tomography shown to outperform conventional angiography for stent guidance in complex cases

September 2 2024

---



Credit: Pixabay/CC0 Public Domain

Using optimal coherence tomography (OCT) to guide stent implantation during percutaneous coronary intervention (PCI) in patients with

complex coronary lesions significantly improves survival and reduces adverse cardiovascular events compared to angiography-guided PCI, the most commonly used method, according to late breaking research presented Sept. 2 in a Hot Line Session at this year's [ESC Congress 2024](#)

"Our results show that the use of OCT guidance improves clinical outcomes and enhances safety and effectiveness of coronary stenting, reducing [cardiac death](#), [myocardial infarction](#), stent thrombosis, and the need for revascularization by 38% in a group of very complex patients, which more reflects contemporary interventional cardiology practice," said lead investigator Professor Byeong-Keuk Kim from Yonsei University Severance Hospital, Seoul, Korea.

Patients with [coronary artery disease](#)—plaque buildup inside the arteries that leads to chest pain, shortness of breath, and [heart attack](#)—often undergo PcCI, a non-surgical procedure that uses a catheter to place stents in the blocked coronary arteries to restore [blood flow](#).

Angiography is the most commonly used method to guide PCI, which involves a special contrast dye and X-ray to see how blood flows through the heart arteries to highlight any blockages.

OCT uses light instead of X-ray to create cross-sectional and longitudinal images, which have much higher resolution and are more accurate and more detailed compared to angiography.

Previous studies have suggested practical benefits of OCT-guided PCI by providing comprehensive information on coronary vessels, plaques, and implanted stents. This latest study follows up on findings of the OCTOBER trial published in 2023 showing that OCT-guided PCI of complex bifurcation lesions is superior to angiography-guided PCI for cardiovascular outcomes at two years.

The OCCUPI trial enrolled 1,604 patients (aged 19 or older) from 20 centers in The Republic of Korea with anatomically complex lesions requiring PCI using drug-eluting stents.

Patients were randomized in a 1:1 ratio to undergo either OCT guidance (803 patients) or angiography guidance (801 patients). In total, 1,588 (99%) patients completed the 1-year follow-up.

The main (primary) measure of interest was a composite of cardiac death, myocardial infarction, stent thrombosis, and target-vessel revascularization (procedure that restores blood flow) at one year.

At one year follow-up, patients who received OCT guidance experienced a 38% reduction in the combined risk of cardiac death, heart attacks, [stent thrombosis](#), and the need for revascularization compared with angiography guidance (37 patients, 4.6% vs. 59 patients, 7.4%, respectively).

Notably, patients who received OCT guidance were 64% less likely to experience a spontaneous myocardial infarction (excluding periprocedural myocardial infarction) or require target-vessel revascularization than those in the angiography-guided group (0.9% vs. 2.4% and 1.5% vs. 4.1%, respectively).

The study also found that the incidence of contrast-induced nephropathy (damage to kidney) after PCI was not significantly different between the two groups (1.3% vs. 0.9%).

"Our findings provide more evidence that OCT guidance of PCI procedures in patients with complex lesions has a greater impact on improving their lives than conventional angiography [guidance](#)," said Professor Kim. "We now need to establish a detailed standard for optimal use of OCT for the improvement of clinical outcomes of PCI

for complex cases based on the OCCUPI trial."

Provided by European Society of Cardiology

Citation: Optical coherence tomography shown to outperform conventional angiography for stent guidance in complex cases (2024, September 2) retrieved 4 September 2024 from

<https://medicalxpress.com/news/2024-09-optical-coherence-tomography-shown-outperform.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.