


# Stimulating specific neurons in the striatum stops compulsive behavior in mice

September 10 2024

---



Interneurons. Credit: UCLA Broad Stem Cell Research Center.

What if we could resist compulsions? These irrational behaviors, particularly common in obsessive-compulsive disorder (OCD), are hard to suppress.

At Paris Brain Institute, Éric Burguière's team shows that we can anticipate and block them before they occur. These promising results, obtained in mice, could help us understand the biological mechanisms underlying compulsions. They [are published](#) in the journal *Nature Neuroscience*.

Repeatedly touching a particular object, biting your nails, washing your hands every ten minutes... these irrepressible rituals sometimes punctuate our lives, especially during periods of high anxiety. When they become invasive to the point of interfering with daily activities, they can become pathological. This is called compulsions—which are characteristic of several mental illnesses, including [obsessive-compulsive disorder](#) (OCD).

Affected patients know that these behaviors are the product of their mental activity, but it does not always help them regain control. The [compulsion](#) sometimes imposes itself so violently that it disrupts professional activity and social life—to the point that patients isolate themselves to accommodate the symptoms.

Recent research has shown that compulsive episodes are associated with dysfunction of brain circuits located in the orbitofrontal cortex—a region involved in decision-making—and in the striatum, beneath the cortex. However, the underlying neuronal anomalies remained to be identified.

"We had a suspect in mind: inhibitory interneurons, a type of nerve cell found in the striatum. Therefore, our team investigated to better understand the role of these cells in regulating behavior," explains Burguière (CNRS), head of the Neurophysiology of Repetitive Behaviors' team at Paris Brain Institute.

## **Striatal interneurons under fire**

First, Lizbeth Mondragón-González and Christiane Schreiweis, in Burguière's team, selected genetically modified mice that are usually used to study OCD because they display a compulsive behavior which is easy to observe and measure: they groom themselves excessively to the point of causing significant hair loss and skin irritation.

The researchers then used an optogenetics technique, which allows for selective activation or inhibition of neurons using light rays to correct faulty neural circuits.

Their results are unequivocal: thanks to the continuous stimulation of the inhibitory interneurons in the striatum, the grooming of the mice became perfectly normal.

"This confirms the involvement of inhibitory neurons in the compulsions of these animals. However, we did not want to stop there! Our goal is to understand when and how to intervene to stop the symptoms," says Burguière.

## **Nipping compulsion in the bud**

The team then used an [artificial intelligence](#) (AI) technique that analyzes [brain activity](#) to extract information that usually escapes the human eye. In this case, AI helped the researchers to pinpoint the warning signs of a compulsive episode in the [orbitofrontal cortex](#): an increase in slow brain waves, known as "Delta" waves.

Thanks to this signature of the coming crisis, the researchers developed a closed-loop neurostimulation system, which consists of briefly exciting the [inhibitory interneurons](#) upstream of the compulsions. The technique was a success: it allowed suppressing compulsive grooming in mice while reducing the number and duration of stimulations.

These advances could thus contribute to developing new neurostimulation therapies—which would be less energy-intensive than current techniques.

"Deep brain stimulation, which involves implanting electrodes to deliver a low-intensity electrical current in the brain, has already proved effective in treating severe obsessive-compulsive disorder," explains Burguière.

"One of the obstacles to this technique is the need to replace the neurostimulator battery regularly. Our study opens the way to approaches that could reduce the need for surgery and make the treatment lighter and more personalized for the patient."

**More information:** Sirenia Lizbeth Mondragón-González et al, Closed-loop recruitment of striatal interneurons prevents compulsive-like grooming behaviors, *Nature Neuroscience* (2024). [DOI: 10.1038/s41593-024-01633-3](https://doi.org/10.1038/s41593-024-01633-3).

Provided by Paris Brain Institute

Citation: Stimulating specific neurons in the striatum stops compulsive behavior in mice (2024, September 10) retrieved 10 September 2024 from <https://medicalxpress.com/news/2024-09-specific-neurons-striatum-compulsive-behavior.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.