


# Soda warning? New study supports link between diabetes, high-fructose corn syrup

August 23 2007

---



New evidence suggests that sodas sweetened with high-fructose corn syrup may increase the risk of diabetes, particularly in children. Credit: American Chemical Society

Researchers have found new evidence that soft drinks sweetened with high-fructose corn syrup (HFCS) may contribute to the development of diabetes, particularly in children. In a laboratory study of commonly consumed carbonated beverages, the scientists found that drinks containing the syrup had high levels of reactive compounds that have been shown by others to have the potential to trigger cell and tissue damage that could cause the disease, which is at epidemic levels. They

reported results at the 234<sup>th</sup> national meeting of the American Chemical Society.

HFCS is a sweetener found in many foods and beverages, including non-diet soda pop, baked goods, and condiments. It has become the sweetener of choice for many food manufacturers because it is considered more economical, sweeter and more easy to blend into beverages than table sugar. Some researchers have suggested that high-fructose corn syrup may contribute to an increased risk of diabetes as well as obesity, a claim which the food industry disputes. Until now, little laboratory evidence has been available on the topic.

In the current study, Chi-Tang Ho, Ph.D., conducted chemical tests among 11 different carbonated soft drinks containing HFCS. He found ‘astonishingly high’ levels of reactive carbonyls in those beverages. These undesirable and highly-reactive compounds associated with “unbound” fructose and glucose molecules are believed to cause tissue damage, says Ho, a professor of food science at Rutgers University in New Brunswick, N.J. By contrast, reactive carbonyls are not present in table sugar, whose fructose and glucose components are “bound” and chemically stable, the researcher notes.

Reactive carbonyls also are elevated in the blood of individuals with diabetes and linked to the complications of that disease. Based on the study data, Ho estimates that a single can of soda contains about five times the concentration of reactive carbonyls than the concentration found in the blood of an adult person with diabetes.

Ho and his associates also found that adding tea components to drinks containing HFCS may help lower the levels of reactive carbonyls. The scientists found that adding epigallocatechin gallate (EGCG), a compound in tea, significantly reduced the levels of reactive carbonyl species in a dose-dependent manner when added to the carbonated soft

drinks studied. In some cases, the levels of reactive carbonyls were reduced by half, the researchers say.

“People consume too much high-fructose corn syrup in this country,” says Ho. “It’s in way too many food and drink products and there’s growing evidence that it’s bad for you.” The tea-derived supplement provides a promising way to counter its potentially toxic effects, especially in children who consume a lot of carbonated beverages, he says.

But eliminating or reducing consumption of HFCS is preferable, the researchers note. They are currently exploring the chemical mechanisms by which tea appears to neutralize the reactivity of the syrup.

Ho’s group is also probing the mechanisms by which carbonation increases the amount of reactive carbonyls formed in sodas containing HFCS. They note that non-carbonated fruit juices containing HFCS have one-third the amount of reactive carbonyl species found in carbonated sodas with HFCS, while non-carbonated tea beverages containing high-fructose corn syrup, which already contain EGCG, have only about one-sixth the levels of carbonyls found in regular soda.

In the future, food and drink manufacturers could reduce concerns about HFCS by adding more EGCG, using less HFCS, or replacing the syrup with alternatives such as regular table sugar, Ho and his associates say.

Source: American Chemical Society

Citation: Soda warning? New study supports link between diabetes, high-fructose corn syrup (2007, August 23) retrieved 20 September 2024 from <https://medicalxpress.com/news/2007-08-soda-link-diabetes-high-fructose-corn.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.