

Sleep disorders: A warning sign for neurodegenerative disease?

December 24 2008

According to the latest study by Dr. Ronald Postuma from the Research Institute of the MUHC and Dr. Jacques Montplaisir from the Université de Montréal and the Hôpital du Sacré-Cœur de Montréal, 52.4 per cent of patients with REM sleep behaviour disorder develop a neurodegenerative disease within 12 years following their initial diagnosis. These results will be published on December 24, 2008 in the journal *Neurology*, the official publication of the American Academy of Neurology.

The study showed that the chance a patient suffering from an REM sleep behaviour disorder will develop a neurodegenerative disease is 17.7 per cent within five years of diagnosis, 40.6 per cent within 10 years, and 52.4 per cent within 12 years. "These results establish a clear link and indicate that these sleep disorders could be a predictor of neurodegenerative disease," explained Dr. Postuma.

The 93 patients who participated in this study were recruited and assessed at the Hôpital du Sacré-Coeur de Montréal by Dr Jacques Montplaisir.

"Doctors should pay close attention when following these patients, as their observations could help define the precursors of diseases such as Parkinson's, Alzheimer's, Lewy body dementia, or multiple system atrophy," stated Dr. Montplaisir, principal investigator of the study. Currently, it is difficult to diagnose these diseases with certainty before an advanced stage, as doctors lack data on warning signs. Understanding

how to detect these diseases early would be of great value to clinical practice.

Although effective treatments against REM sleep behaviour disorder do exist, these medications do not postpone the onset of neurodegenerative disease. As research is very active in this field, these patients could represent a viable target population in the relatively near future to test the effectiveness of new innovative treatments to fight neuronal degeneration.

REM sleep behaviour disorder affects a small percentage of the population. It is characterized by a loss of the normal muscle relaxation while dreaming and is seen most often in men fifty and older. This is a specific pathology that should not be confused with insomnia, night terrors, or confusional arousals.

Source: McGill University Health Centre

Citation: Sleep disorders: A warning sign for neurodegenerative disease? (2008, December 24) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2008-12-disorders-neurodegenerative-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--