

# Liver cell adenoma or hepatocellular carcinoma?

March 20 2009

---

Liver cell adenoma (LCA) is a benign tumor of the liver parenchyma that is associated with the use of oral contraceptives or with glycogen-storage disease. A group in Japan reported a case of LCA found in a 40-year-old woman without a history of oral contraceptive use in which the sequential alteration of the radiological findings suggested well-differentiated hepatocellular carcinoma.

Recently, LCAs with typical backgrounds of the patients are diagnosed by radiological findings without pathological findings due to the progress of diagnostic imaging techniques. A typical LCA shows low density or isodensity on plain computed tomography, presents a homogenous contrasting effect on arterial phase and does not show an apparent wash out on delayed phase.

A typical LCA in magnetic resonance imaging shows almost the same signal intensity as the surrounding parenchyma on T1 and T2 and shows high intensity on fat suppression T2. However, LCA often presents difficulties in a differential diagnosis with well-differentiated HCC, especially without the typical backgrounds of the patients.

A research article to be published on March 14, 2009 in the *World Journal of Gastroenterology* reports an extremely rare case of LCA. The research team led by Prof. Shimosegawa from the Division of Gastroenterology, Tohoku University reported a case of LCA found in a 40-year-old woman without a typical history in which the sequential alteration of the radiological findings suggested well-differentiated

HCC.

A liver tumor 35 mm in diameter was found incidentally in a woman, which showed a homogenous enhancement in the arterial phase and almost the same enhancement as the surrounding [liver parenchyma](#) in the delayed phase of CT. The tumor was found to contain fat on MRI. However, radiological findings altered, which caused to suspect that a well-differentiated HCC containing fat was becoming dedifferentiated. Partial hepatectomy was performed and the pathological findings showed the typical findings of LCA. Diagnosis of LCA only by radiological findings is difficult in patients without backgrounds such as a history of receiving contraceptives or glycogen-storage disease.

More information: Kogure T, Ueno Y, Sekiguchi S, Ishida K, Igarashi T, Wakui Y, Iwasaki T, Shimosegawa T. [Liver cell](#) adenoma showing sequential alteration of radiological findings suggestive of well-differentiated hepatocellular carcinoma. World J Gastroenterol 2009; 15(10): 1267-1272 [www.wjgnet.com/1007-9327/15/1267.asp](http://www.wjgnet.com/1007-9327/15/1267.asp)

Source: World Journal of Gastroenterology

Citation: Liver cell adenoma or hepatocellular carcinoma? (2009, March 20) retrieved 19 September 2024 from <https://medicalxpress.com/news/2009-03-liver-cell-adenoma-hepatocellular-carcinoma.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--