

# Avoiding hysterectomy: Major interventional radiology E-collection info available

14 July 2009

For the first time, the Society of Interventional Radiology has assembled a major electronic collection of professional articles about uterine artery embolization, a treatment directed toward a number of conditions involving the uterus—most often adverse health effects that may occur due to the presence of uterine fibroids. The *Journal of Vascular and Interventional Radiology* "virtual" collection allows health care providers and the public to view the abstracts on current research on this topic in one place, eliminating the need to search topics individually.

Uterine fibroids are very common noncancerous growths that develop in the muscular wall of the [uterus](#). Hysterectomy, surgical removal of the uterus performed by a gynecologist, is the most common treatment for symptomatic uterine fibroids. In fact, more than 200,000 women in the United States have hysterectomies for fibroids each year; it is the second most common surgery among women. However, most women are candidates for uterine artery embolization (also called uterine fibroid embolization). "Uterine artery embolization is a treatment method for fibroids that is relatively noninvasive and has high success rates. Women with fibroids and their health care providers should be aware of this therapy. This collection assembles relevant information about the interventional radiology treatment in one convenient place," explained Albert A. Nemcek Jr., M.D., FSIR, editor of *JVIR*, a peer-reviewed, monthly publication long recognized for its exceptional quality and influence as an academic and professional resource.

"The value for physicians and patients is the convenience of having all the recent articles from a major journal in one place. While it is not a comprehensive list of all recent uterine artery embolization articles, many of the major studies are included, particularly those that relate to the technical aspects of the procedure," said James B. Spies, M.D., FSIR, *JVIR* section editor for women's

health. "Nearly all the new innovations in the procedure have been reported in *JVIR* first. The collection has particular value to physicians who are just entering the field or who are in training, as it provides a continuum of the recent innovations in the procedure and is a good example of how medical research progresses over time," added Spies, professor of radiology and chair of the Department of Radiology at Georgetown University Medical Center in Washington, D.C.

Uterine artery embolization, a minimally invasive, targeted treatment, offers less risk, less pain and less recovery time than traditional surgery. Interventional radiologists use expertise in imaging to deliver treatment directly to a fibroid, blocking the flow of blood to the tumor and causing it to shrink. "With these modern treatments, hopefully the trend to offer traditional surgery first will be reversed. It's important for patients to ask questions, obtain consults with different types of doctors and know all their treatment options," added Nemcek, an interventional radiologist and professor of radiology and surgery at Northwestern Memorial Hospital in Chicago, Ill.

The 46-article collection of previously published articles and abstracts—with some of the leading researchers in the field—is divided into nine sections: indications, uterine imaging, technique, outcomes, other indications for embolization, economics, radiation exposure, other procedures and animal studies.

According to Spies, the collection includes an important article, "Leiomyoma Infarction After Uterine Artery Embolization: A Prospective Randomized Study Comparing Tris-acryl Gelatin Microspheres Versus Polyvinyl Alcohol Microspheres," a randomized study by Gary P. Siskin, M.D., FSIR, and colleagues at Albany Medical College, that answers a key technical question regarding choice of embolic material. Also important are "Economic Evaluation of Uterine

Artery Embolization Versus Hysterectomy in the Treatment of Symptomatic Uterine Fibroids: Results From the Randomized EMMY Trial" and "Payer Costs in Patients Undergoing Uterine Artery Embolization, [Hysterectomy](#) or Myomectomy for [Treatment](#) of Uterine Fibroids." These latter two articles provide strong evidence—obtained in two different ways—that [uterine artery](#) embolization is less costly than traditional surgery, said Spies.

Source: Society of Interventional Radiology ([news](#) : [web](#))

APA citation: Avoiding hysterectomy: Major interventional radiology E-collection info available (2009, July 14) retrieved 17 May 2021 from <https://medicalxpress.com/news/2009-07-hysterectomy-major-interventional-radiology-e-collection.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*