

Researchers identify gene that spurs deadly brain cancer

December 3 2009

Howard Hughes Medical Institute (HHMI) researchers have identified a new factor that is necessary for the development of many forms of medulloblastoma, the most common type of malignant childhood brain cancer.

HHMI investigator Huda Y. Zoghbi and colleagues at Baylor College of Medicine prevented medulloblastoma from developing in mice by shutting down production of the protein Atoh1 in susceptible [brain cells](#). The team's findings, reported in the December 4, 2009, issue of *Science*, suggest Atoh1 may be a new target for medulloblastoma treatment.

"When we cloned the gene for Atoh1 in 1996, we had no clue that it had any medical relevance," said Zoghbi, a neuroscientist and neurologist. "Now we know that it's critical for many medical issues, the most recent one being this common childhood cancer."

Atoh1 (also known as Math1) is a transcription factor that works in the nuclei of cells to keep certain genes switched on. It is evolutionarily ancient, appearing in slightly varying forms in various species, from [fruit flies](#) to humans. In cells where Atoh1 is active, it seems to be switched on only during [fetal development](#), when cells proliferate rapidly to fill out the various parts of the [nervous system](#).

However, in the region of the [brain](#) known as the cerebellum, Atoh1 is active after birth in the fast-dividing granule neuron precursor (GNPs) cells that eventually stop dividing and become mature granule neurons.

"The cerebellar granule neurons are unique in that most of their development happens after birth, both in mice and humans," Zoghbi said.

A few years ago, experiments done in several laboratories hinted that Atoh1 might be required to keep GNPs in their fast-dividing state and make them more susceptible to developing into medulloblastoma tumors.

"The question for us was whether we could really prove, not just in the cell culture dish or in microarrays but in animals, that Atoh1 plays this role in medulloblastoma," Zoghbi said.

Ordinarily, to begin to discern the function of a gene such as Atoh1, researchers would engineer a strain of mice that lack the gene. But that had been tried in the 1990s, and the results were less than satisfying. Researchers found that Atoh1-knockout mice failed to develop properly in the womb, and died at birth. To study Atoh1's function after birth, Zoghbi's team, led by postdoctoral researcher Adriano Flora, devised a more advanced technique. First they bred a strain of mice with a genetic off-switch connected to their Atoh1 gene; then they injected a chemical into the brains of healthy newborn mice, to trigger this off-switch and eliminate the production of Atoh1 in GNPs. As a result, the GNPs immediately stopped proliferating and started maturing into granule neurons.

That result showed that Atoh1 helped keep GNPs in their ever-dividing state. Further experiments revealed that Atoh1 revs up GNPs by switching on a gene called Gli2, a well-known member of the Sonic Hedgehog signaling pathway that helps cells divide. The Sonic Hedgehog pathway is also inappropriately switched on in many cancers, including medulloblastoma.

"At this point we asked whether we could affect the development of

medulloblastoma in mice by shutting down Atoh1," Zoghbi said.

To find out, the team applied their local Atoh1-shutdown technique to a special strain of mice with a specific genetic mutation that makes them develop medulloblastoma. In these [mice](#), a mutant gene is switched on after birth, sending the Sonic Hedgehog signaling pathway into overdrive, causing precancerous lesions and tumors in the cerebellum. But when Zoghbi's team switched off Atoh1, these cancerous changes never occurred.

Establishing Atoh1 as a key player in the origin of medulloblastoma makes it a potential target for new drug treatments, Zoghbi said. But to Zoghbi, an important next step is to determine whether the protein is still needed to keep tumors growing after they've become established: "If we allow these tumors to develop, and then we take away Atoh1, would that make a difference?" Her lab and others are also now racing to determine what keeps Atoh1 inappropriately switched on in medulloblastoma cells, and what normally switches it off.

Zoghbi emphasized that she originally took up the study of Atoh1 as an exercise in pure biology, with no idea that it would have relevance to disease. "That just underscores the tremendous importance of doing science for science's sake," she said.

Source: Howard Hughes Medical Institute ([news](#) : [web](#))

Citation: Researchers identify gene that spurs deadly brain cancer (2009, December 3) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2009-12-gene-spurs-deadly-brain-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.