

Diet May Protect Against Gene Changes in Smokers

12 January 2010

(PhysOrg.com) -- Leafy green vegetables, folate, and some multivitamins could serve as protective factors against lung cancer in current and former smokers, according to a study that is a first step in understanding a complex association. The study was supported by the National Cancer Institute (NCI), part of the National Institutes of Health. The study appeared online Jan. 12, 2010, in *Cancer Research*.

Researchers, led by Steven Belinsky, Ph.D., Lovelace Respiratory Research Institute, Albuquerque, N.M., examined cells that were coughed up by current and former smokers. Upon careful study of the cells and by comparing those cells with profiles of smokers' dietary intake of leafy green [vegetables](#), folate, and some multivitamins, they found that those particular substances could influence the prevalence of cellular gene methylation. Gene methylation is a chemical modification used by the cell to control [gene expression](#).

The addition of methyl groups, which are simple four-atom molecules, to DNA can affect whether the gene is expressed, i.e., whether the gene's signal to produce a protein is actually sent. Many [genes](#) involved in critical cell functions, including cell division, are methylated in lung tumors. As seen in previous studies, gene methylation is likely to be a major mechanism in lung [cancer development](#) and progression, as well as a potential marker for the early detection of [lung cancer](#).

In the study, slightly more than 1,100 current and former smokers from the Lovelace Smokers Cohort submitted sputum samples and completed questionnaires regarding their dietary intake. Most (75 percent) of the participants were women who had been enrolled in the study since 2001. Men were not enrolled until 2004.

The researchers analyzed cells in the sputum

samples for the methylation status of eight genes that were linked to increased risk for lung cancer in previous studies. Two of the genes, p16 and MGMT, have been found to be highly associated with increased methylation rates.

The researchers next investigated associations between 21 dietary variables and methylation. Both higher intake of leafy green vegetables and higher intake of folate were significantly associated with a reduced probability of high methylation. Current multivitamin users also had less DNA methylation, although there was no association between the duration of multivitamin use and methylation.

"Aberrant gene methylation is a known mechanism in the development of cancer from cigarette smoke carcinogens," said Jacob Kagan, Ph.D., of the Cancer Biomarkers Research Group in NCI's Division of Cancer Prevention.

There has been considerable debate regarding the relationship between diet and cancer prevention. Previous studies, such as the Alpha-Tocopherol, Beta-Carotene Cancer Prevention (ATBC) Trial, showed an increased risk of lung cancer in smokers who took beta-carotene supplements. In contrast, the current study shows reduced gene methylation with the intake of multivitamin supplements that are rich in phytochemicals, such as vitamin C, carotenoids, lutein, folic acid, and vitamins A and K — the same agents present in leafy greens.

Previous studies have suggested an association between a low folate intake and increased lung cancer risk in current and former [smokers](#). Higher folate intake has been associated with lower methylation of genes in colorectal tumors as well.

"Additional research is needed to independently validate the current observations, and also to help resolve contradictions between varying studies. This particular study used a well-planned design

and can serve as a basis for future identification of the mechanistic targets of these dietary factors. Such studies are important steps for the future success of chemopreventive strategies" said Sudhir Srivastava, Ph.D., chief, Biomarkers Research Group, NCI.

The authors of this study believe that since gene methylation is a promising marker for lung cancer, understanding the factors underlying methylation is a high priority and could be used for early detection and chemoprevention of lung cancer. They also recommend that lung cancer prevention interventions be developed that take into consideration the influence of dietary factors on cancer risk.

More information: Stidley CA, Picchi MA, Leng S, Willink R, Crowell R, Flores KG, Kang H, Byers T, Gilliland FD, and Belinsky SA. Multi-Vitamins, Folate, and Green Vegetables Protect Against Gene Promoter Methylation in the Aerodigestive Tract of Smokers. Online Jan. 12, 2010, Cancer Research.

Provided by National Institutes of Health

APA citation: Diet May Protect Against Gene Changes in Smokers (2010, January 12) retrieved 7 December 2021 from <https://medicalxpress.com/news/2010-01-diet-gene-smokers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.