

Rectal cancer rates are rising in young individuals

23 August 2010

A new analysis has found that while colon cancer rates have remained steady over the past several decades among people under the age of 40, rectal cancer rates are increasing in this population across races and in both sexes. Published early online in *Cancer*, a peer-reviewed journal of the American Cancer Society, the study indicates that greater efforts are needed to diagnose rectal cancer in young individuals who show potential signs of the disease.

Rectal cancer is considered to be rare among young individuals in the United States. Because underestimating rectal cancer's incidence may lead to missed or delayed diagnoses in younger people, Joshua Meyer, MD, a radiation oncologist currently at Fox Chase Cancer Center, led a team that analyzed trends in rectal cancer incidence in the United States compared with [colon cancer](#) trends. Dr. Meyer worked on this research while at the New York-Presbyterian Hospital/Weill Cornell Medical Center in New York City.

By conducting a retrospective study using data from the Surveillance Epidemiology and End Results (SEER) cancer registry, the investigators identified 7,661 colon and rectal cancer patients under age 40 years between 1973 and 2005. The researchers then calculated the change in incidence over time for colon and rectal cancers.

Overall rates of colon and of rectal cancer were low during the years of the study (1.11 cases and 0.42 cases per 100,000, respectively). While colon cancer rates remained essentially flat in individuals under age 40 years in recent decades, [rectal cancer rates](#) have been increasing since 1984. Specifically, between 1984 and 2005, the rate of rectal cancer diagnosis rose 3.8% per year.

"We suggest that in young people presenting with rectal bleeding or other common signs of rectal cancer, endoscopic evaluation should be considered in order to rule out a malignancy," said

Dr. Meyer. "This is in contrast to what is frequently done, which is to attribute these findings to hemorrhoids. More frequent endoscopic evaluation may be able to decrease the documented delay in diagnosis among young people," he explained. Because the overall incidence of [rectal cancer](#) is relatively low, the authors do not advocate for a change in screening guidelines.

More information: "Increasing incidence of rectal cancer in patients under age 40: an analysis of the Surveillance, Epidemiology, and End Results database," Joshua E. Meyer, Tarun Narang, Felice H. Schnoll-Sussman, Mark B. Pochapin, Paul J. Christos, David L. Sherr. *Cancer*; Published Online: August 23, 2010 ([DOI: 10.1002/cncr.25432](https://doi.org/10.1002/cncr.25432)).

Provided by American Cancer Society

APA citation: Rectal cancer rates are rising in young individuals (2010, August 23) retrieved 19 November 2019 from <https://medicalxpress.com/news/2010-08-rectal-cancer-young-individuals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.