

Secondhand television exposure linked to eating disorders

January 6 2011

For parents wanting to reduce the negative influence of TV on their children, the first step is normally to switch off the television set.

But a new study suggests that might not be enough. It turns out indirect media exposure, i.e., having friends who watch a lot of TV, might be even more damaging to a teenager's body image.

Researchers from Harvard Medical School's Department of Global Health and Social Medicine examined the link between media consumption and eating disorders among adolescent girls in Fiji.

What they found was surprising. The study's subjects did not even need to have a [television](#) at home to see raised risk levels of eating disorder symptoms.

In fact, by far the biggest factor for eating disorders was how many of a subject's friends and schoolmates had access to TV. By contrast, researchers found that direct forms of exposure, like personal or parental viewing, did not have an independent impact, when factors like urban location, [body shape](#) and other influences were taken into account.

It appeared that changing attitudes within a group that had been exposed to television were a more powerful factor than actually watching the programs themselves. In fact, higher peer [media exposure](#) were linked to a 60 percent increase in a girl's odds of having a high level of eating disorder symptoms, independently of her own viewing.

Lead author Anne Becker, vice chair of the Department of Global Health and Social Medicine at Harvard Medical School, said this was the first study to attempt to quantify the role of social networks in spreading the negative consequences of [media consumption](#) on eating disorders.

"Our findings suggest that social network exposure is not just a minor influence on eating pathology here, but rather, IS the exposure of concern," she said.

"If you are a parent and you are concerned about limiting cultural exposure, it simply isn't going to be enough to switch off the TV. If you are going to think about interventions, it would have to be at a community or peer-based level."

Becker hopes the paper will encourage debate about responsible programming and the regulation of media content to prevent children from secondhand exposure.

"Up until now, it has been very difficult to get people who produce media as entertainment to come to the table and think about how they might ensure that their products are not harmful to children," she said.

This is Becker's second study of media's impact in Fiji, which is an ideal location for broadcast media research because of the recent arrival of television, in the 1990s, and the significant regional variations in exposure to TV, the Internet and print media. Some remote areas in the recent study still did not have electricity, cell phone reception, television or the Internet when the data were collected in 2007.

Her first study found a rise in [eating disorder](#) symptoms among [adolescent girls](#) following the introduction of broadcast television to the island nation in 1995.

What makes Fiji a particularly interesting case is that traditional culture prizes a robust body shape, in sharp contrast to the image presented by Western television shows such as Beverly Hills 90210, Seinfeld and Melrose Place, which were quite popular in Fiji when television debuted there in the 1990s.

Girls would see actresses as role models, says Becker, and began noting how a slender body shape was often accompanied by success in those shows. This perception appears to have been one of the factors leading to a rise in eating pathology among the Fijian teenagers.

But until now, it was not known how much of this effect came from an individual's social network.

Nicholas Christakis, professor of medical sociology in the department of health care policy at Harvard Medical School, has studied the spread of health problems through social networks.

"It shouldn't be that surprising to us, even though it is intriguing, that the indirect effects of media are greater," Christakis said. "Most people aren't paying attention to the media, but they are paying attention to what their friends say about what's in the media. It's a kind of filtration process that takes place by virtue of our social networks."

Becker says that although the study focused on Fijian schoolgirls, remote from the US, it warrants concern and further investigation of the health impact on other populations.

More information: "Social network media exposure and adolescent eating pathology in Fiji" *The British Journal of Psychiatry* (2011), 198, 43-50

Provided by Harvard Medical School

Citation: Secondhand television exposure linked to eating disorders (2011, January 6) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2011-01-secondhand-television-exposure-linked-disorders.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.