

Dopamine controls formation of new brain cells

8 April 2011

(PhysOrg.com) -- A study of the salamander brain has led researchers at Karolinska Institutet to discover a hitherto unknown function of the neurotransmitter dopamine. In an article published in the prestigious scientific journal *Cell Stem Cell* they show how in acting as a kind of switch for stem cells, dopamine controls the formation of new neurons in the adult brain. Their findings may one day contribute to new treatments for neurodegenerative diseases, such as Parkinson's.

The study was conducted using salamanders which unlike mammals recover fully from a Parkinson's-like condition within a four week period. Parkinson's disease is a neurodegenerative disease characterised by the death of dopamine-producing cells in the mid-brain. As the salamander re-builds all lost dopamine-producing neurons, the researchers examined how the salamander [brain](#) detects the absence of these cells. This question is a fundamental one since it has not been known what causes the new formation of nerve cells and why the process ceases when the correct number have been made.

What they found out was that the salamander's [stem cells](#) are automatically activated when the [dopamine](#) concentration drops as a result of the death of dopamine-producing neurons, meaning that the neurotransmitter acts as a constant handbrake on stem cell activity.

"The medicine often given to Parkinson's patients is L-dopa, which is converted into dopamine in the brain," says Dr Andras Simon, who led the study at the Department of Cell and Molecular Biology. "When the [salamanders](#) were treated with L-dopa, the production of new dopamine-producing neurons was almost completely inhibited and the animals were unable to recover. However, the converse also applies. If dopamine signalling is blocked, new neurons are born unnecessarily."

As in mammals, the formation of neurons in the

salamander mid-brain is virtually non-existent under normal circumstances. Therefore by studying the salamander, scientists can understand how the production of new nerve cells can be resumed once it has stopped, and how it can be stopped when no more [neurons](#) are needed. It is precisely in this regulation that dopamine seems to play a vital part. Many observations also suggest that similar mechanisms are active in other animal species too. Further comparative studies can shed light on how neurotransmitters control stem cells in the brain, knowledge that is of potential use in the development of therapies for [neurodegenerative diseases](#).

"One way of trying to repair the brain in the future is to stimulate the stem cells that exist there," says Dr Simon. "This is one of the perspectives from which our study is interesting and further work ought to be done on whether L-dopa, which is currently used in the treatment of Parkinson's, could prevent such a process in other species, including humans. Another perspective is how medicines that block dopamine signalling and that are used for other diseases, such as psychoses, affect stem cell dynamics in the brain."

The salamander is a tailed member of the frog family most known for its ability to regenerate lost body parts, such entire limbs.

More information: Daniel A Berg, Matthew Kirkham, Heng Wang, Jonas Frisén & Andras Simon, Dopamine Controls Neurogenesis in the Adult Salamander Midbrain in Homeostasis and during Regeneration of Dopamine Neurons, *Cell Stem Cell*, online 7 April 2011. www.cell.com/cell-stem-cell/

Provided by Karolinska Institutet

APA citation: Dopamine controls formation of new brain cells (2011, April 8) retrieved 7 December 2021 from <https://medicalxpress.com/news/2011-04-dopamine-formation-brain-cells.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.