

Rehab robots lend stroke patients a hand

August 11 2011

Robot-assisted therapy has measurable benefits for patients with a weaker arm following a stroke. This is according to new research featured in the journal *Clinical Rehabilitation*, published by SAGE, which is the first to use accelerometers to track patients' improvement and compare real world results.

The study authors, Keh-chung Lin, Yu-wei Hsieh, Wan-wen Liao - National Taiwan University, Ching-yi Wu - Chang Gung University, and Wan-ying Chang, Department of Physical Medicine and Rehabilitation, Taipei Hospital aimed to investigate how robot-assisted [therapy](#) helps [arm function](#) to improve after a stroke. They enrolled 20 patients in a study comparing robot-assisted therapy combined with functional training against an active control treatment group.

[Stroke patients](#) usually have difficulties transferring motor skills learned in therapy to their daily living environment because of cognitive deficits. The authors included real-world arm activity in the study by having patients wear accelerometers on both arms daily as they went about their normal tasks.

One of the key findings of the study was that robot-assisted therapy, when combined with functional task training, helps functional arm use and improves bimanual arm activity in daily life. Patients following a stroke often have weakness on one side of the upper body (hemiparesis), which can make daily life more difficult. Robotic rehabilitation is increasingly available, and holds promise for enhancing traditional post-stroke interventions. Because robots never tire, they can provide massive

and intensive training in a consistent manner without fatigue, with programming precisely tailored to each patient's needs.

The rehab robots give sensorimotor feedback via visual and [auditory feedback](#) during training sessions, to facilitate patients' motor learning. However, although stroke patients' arm motor function and [muscle strength](#) have shown to improve during robot-assisted therapy in rehabilitation, previous studies suggested that these improvements did not carry through to the patients' daily lives. Some reasons for this might include a need for better measurement scales for patients' real life daily functions, as well as the fact many people compensate by using their non-impaired arm instead. By measuring both arms and following patients with the accelerometers at home, this study addressed these issues.

Two further important findings of the study were that accelerometers are suitable tools to measure real-world arm activity in stroke patients; and that when combined with traditional clinical measurements these can enhance holistic understanding of a patients' life performance.

Accelerometers provide objective information about physical activity by measuring the acceleration of body movements. Stroke patients can easily wear the portable [accelerometers](#) like a wristwatch on each arm. This means researchers now have the accurate information they need to verify the intensity and amount of physical activity the patients actually do in real-life situations.

During the study, both groups received [intensive training](#) for 90 to 105 minutes per session, five days per week for four weeks. The training programs were administered by certified occupational therapists during regularly scheduled training sessions, and all other routine interdisciplinary stroke rehabilitation was continued as usual. Therapy in the control group was designed to match the robot-assisted therapy in amount of therapy hours, and these participants served as a dose-

matched comparison group. Occupational therapists designed activities for the control group based on neurodevelopmental techniques and contemporary rehabilitative models such as task-oriented training and motor learning theory.

The mean ratio change of the robot-assisted therapy group was 0.047 ± 0.047 , which beat the 0.007 ± 0.026 in the control group. The robot-assisted therapy group also handled more daily tasks with their impaired arm than the control group.

"In this study of rehabilitation approaches for [patients](#) with mild-to-moderate upper limb impairment six months after a stroke, we found significantly greater benefits of robot-assisted therapy compared with the active control group on the amount and quality of functional arm activity for the hemiplegic hand in the living environment," said Keh-chung Lin. "Moreover, robot-assisted therapy had superior benefits on improving bimanual arm activity," he added.

Larger studies along with follow up research to look at whether these improvements are lasting will be the next steps towards making the most of robots for stroke patients' rehabilitation in future.

More information: Effects of Robot-Assisted Upper Limb Rehabilitation on Daily Function and Real-World Arm Activity in Patients with Chronic Stroke by Wan-wen Liao, Ching-yi Wu, Yu-wei Hsieh, Keh-chung Lin and Wan-ying Chang published today, August 10th, in *Clinical Rehabilitation*.

Provided by SAGE Publications

Citation: Rehab robots lend stroke patients a hand (2011, August 11) retrieved 24 April 2024

from <https://medicalxpress.com/news/2011-08-rehab-robots-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.