

# Fat cells in abdomen fuel spread of ovarian cancer

30 October 2011

A large pad of fat cells that extends from the stomach and covers the intestines provides nutrients that promote the spread and growth of ovarian cancer, reports a research team based at the University of Chicago in the journal *Nature Medicine*, published online October 30th, 2011.

Ovarian cancer, the fifth leading cause of cancer deaths in women, tends to spread within the [abdominal cavity](#) as opposed to distant organs. In 80 percent of women, by the time ovarian cancer is diagnosed, it has spread to the pad of [fat cells](#), called the omentum. Often, [cancer growth](#) in the omentum exceeds the growth of the original ovarian cancer.

"This [fatty tissue](#), which is extraordinarily rich in energy-dense lipids, acts as a launching pad and energy source for the likely lethal spread of ovarian cancer," said study author Ernst Lengyel, MD, PhD, professor of [obstetrics and gynecology](#) at the University of Chicago. "The cells that make up the omentum contain the biological equivalent of jet fuel. They feed the cancer cells, enabling them to multiply rapidly. Gaining a better understanding of this process could help us learn how to disrupt it."

The researchers performed a series of experiments to identify the role of these fat cells as major mediators of ovarian [cancer metastasis](#). The first step was to understand the biological signals that attract ovarian cancer cells to the omentum and use it for rapid growth.

The spread of ovarian cancer cells to the omentum can happen quickly. Ovarian cancer cells injected into the abdomen of healthy mice find their way to the omentum within 20 minutes. The researchers found that protein signals emitted by the omentum can attract the tumor cells. Inhibitors which disturbed these signals reduced this attraction by at least 50 percent.

Once ovarian cancer cells reach the omentum,

they quickly develop the tools to devour the sustenance provided by this fatty tissue, reprogramming their metabolism to thrive on lipids acquired from fat cells. Ovarian cancer can rapidly convert the entire omentum, a soft fat pad, into a solid mass of cancer cells.

"This mechanism may not be limited to [ovarian cancer cells](#)," the authors note. Fat metabolism may also contribute to cancer development in other environments where fat cells are abundant, such as breast cancer.

A protein known as fatty acid binding protein (FABP4), a fat carrier, may be crucial to this process and could be a target for treatment.

When the researchers compared primary ovarian cancer tissue with [ovarian cancer](#) tissue which had spread to the omentum, they found that [tumor cells](#) next to omental fat cells produced high levels of FABP4. Cancer cells distant from the fat cells did not produce FABP4.

When they inhibited FABP4, the transfer of nutrients from fat cells to cancer cells was drastically reduced. Inhibition of FABP4 also reduced tumor growth and the ability of tumors to generate new blood vessels.

"Therefore," the authors wrote, "FABP4 emerges as an excellent target in the treatment of intra-abdominally disseminating tumors, which preferentially metastasize to adipose tissue such as ovarian, gastric, and colon cancers."

Provided by University of Chicago

APA citation: Fat cells in abdomen fuel spread of ovarian cancer (2011, October 30) retrieved 22 May 2019 from <https://medicalxpress.com/news/2011-10-fat-cells-abdomen-fuel-ovarian.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*