


Brain imaging study finds evidence of basis for caregiving impulse

March 16 2012



MRI brain scan

Distinct patterns of activity-- which may indicate a predisposition to care for infants -- appear in the brains of adults who view an image of an infant face -- even when the child is not theirs, according to a study by researchers at the National Institutes of Health and in Germany, Italy, and Japan.

Seeing images of infant faces appeared to activate in the adult's brains [circuits](#) that reflect preparation for movement and speech as well as [feelings](#) of reward.

The findings raise the possibility that studying this activity will yield insights into care giving behavior, but also in cases of [child neglect](#) or

abuse.

"These adults have no children of their own. Yet images of a baby's face triggered what we think might be a deeply embedded response to reach out and care for that child," said senior author Marc H. Bornstein, Ph.D., head of the Child and Family Research Section of the Eunice Kennedy Shriver National Institute of Child Health and Human Development, the NIH institute that collaborated on the study.

While the researchers recorded participants' [brain](#) activity, the participants did not speak or move. Yet their brain activity was typical of patterns preceding such actions as picking up or talking to an infant, the researchers explained. The activity pattern could represent a biological impulse that governs adults' interactions with small children.

From their study results, the researchers concluded that this pattern is specific to seeing human infants. The pattern did not appear when the participants looked at photos of adults or of animals—even baby animals.

Along with Dr. Bornstein, the research was carried out by first author Andrea Caria, Ph.D., of the University of Tuebingen, in Germany; Paola Venuti of the Department of Cognitive Science of University of Trento in Italy; Gianluca Esposito of the RIKEN Brain Science Institute in Saitama, Japan; researchers from the Max Planck Institute for Biological Cybernetics and Eberhard Karls University, in Tuebingen, Germany.

Their findings appear in the journal *NeuroImage*.

To collect the data, the researchers showed seven men and nine women a series of images while recording their brain activity with a functional magnetic resonance imaging scanner. In the scanner, participants viewed images of puppy and kitten faces, full-grown dogs and cats, human

infants and adults.

When the researchers compared the areas and strength of brain activity in response to each kind of image, they found that infant images evoked more activity than any of the other images in brain areas associated with three main functions:

- Premotor and preverbal activity - The researchers documented increased activity in the premotor cortex and the supplemental motor area, which are regions of the brain directly under the crown of the head. These regions orchestrate brain impulses preceding speech and movement but before movement takes place.
- Facial recognition - Activity in the fusiform gyrus—on each side of the brain, about where the ears are—is associated with processing of information about faces. Activity the researchers detected in the fusiform gyrus may indicate heightened attention to the movement and expressions on an infant's face, the researchers said.
- Emotion and reward - Activity deep in the brain areas known as the insula and the cingulate cortex indicated emotional arousal, empathy, attachment and feelings linked to motivation and reward, the researchers said. Other studies have documented a similar pattern of activity in the brains of parents responding to their own infants.

Participants also rated how they felt when viewing adult and infant faces. They reported feeling more willing to approach, smile at, and communicate with an infant than an adult. They also recorded feeling happier when viewing images of infants.

Taken together, the researchers contend, the findings suggest a readiness to interact with infants that previously has been only inferred, and only

from parents. Such [brain activity](#) in nonparents could indicate that the biological makeup of humans includes a mechanism to ensure that infants survive and receive the care they need to grow and develop.

However, signs of readiness to care for a child that appear in the brains of some or even most adults do not necessarily mean the same patterns will appear in the brains of all adults, Dr. Bornstein said. "It's equally important to investigate what's happening in the brains of those who have neglected or abused children," he said. "Additional studies could help us confirm and understand what appears to be a parenting instinct in [adults](#), both when the instinct functions and when it fails to function."

Provided by National Institutes of Health

Citation: Brain imaging study finds evidence of basis for caregiving impulse (2012, March 16) retrieved 19 September 2024 from <https://medicalxpress.com/news/2012-03-brain-imaging-evidence-basis-caregiving.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--