

Combination of smoking and heavy drinking 'speeds up cognitive decline'

July 10 2013

The combination of smoking and heavy drinking speeds up cognitive decline, according to new research published in the *British Journal of Psychiatry*.

Researchers from UCL (University College London) found that smokers who drank alcohol heavily had a 36% faster cognitive decline compared to non-smoking moderate drinkers.

Smoking and heavier [alcohol consumption](#) often co-occur, and their combined effect on cognition may be larger than the sum of their individual effects. The research team assessed 6,473 adults (4,635 men and 1,838 women) aged between 45 and 69 years old over a 10-year period. The adults were part of the Whitehall II [cohort study](#) of British civil servants.

All the participants were asked about their cigarette and alcohol consumption, and their cognitive function (including verbal and mathematical reasoning, short-term [verbal memory](#) and [verbal fluency](#)) was then assessed three times over 10 years.

The research team found that in current smokers who were also heavy drinkers, cognitive decline was 36% faster than in non-smoking moderate drinkers. This was equivalent to an age effect of 12 years – an additional two years over the 10-year follow up period. Among smokers, cognitive decline was found to be faster as the number of alcohol units consumed increased.

Lead researcher Dr Gareth Hagger-Johnson said: "Our research shows that cognitive decline was 36% faster in those people who reported both cigarette smoking and drinking alcohol above the recommended limits (14 units per week for women, 21 units per week for men). When we looked at people who were heavy-drinking smokers, we found that for every 10 years that they aged their brains aged the equivalent of 12 years."

"From a public health perspective, the increasing burden associated with cognitive aging could be reduced if lifestyle factors can be modified, and we believe that people should not drink alcohol more heavily in the belief that alcohol is a protective factor against cognitive decline. Current advice is that smokers should stop or cut down, and people should avoid heavy alcohol drinking. Our study suggests that people should also be advised not to combine these two unhealthy behaviours—particularly from mid-life onwards. Healthy behaviours in midlife may prevent [cognitive decline](#) into early old age."

More information: Hagger-Johnson G, Sabia S, Brunner EJ, Shipley M, Bobak M, Marmot M, Kivimaki M and Singh-Manoux A. The combined impact of smoking and heavy alcohol use on cognitive decline in early old age: The Whitehall II cohort study. *British Journal of Psychiatry*, ePub ahead of print 11 July 2013

Provided by University College London

Citation: Combination of smoking and heavy drinking 'speeds up cognitive decline' (2013, July 10) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-07-combination-heavy-cognitive-decline.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.