

Study examines risk of severe blood sugar swings among diabetics taking fluoroquinolones

15 August 2013

Diabetic patients taking oral fluoroquinolones, a frequently prescribed class of antibiotics, were found to have a higher risk of severe blood sugar-related problems than diabetic patients taking other kinds of antibiotics, according to a recent study from Taiwan published in *Clinical Infectious Diseases*. The increased risk was low—hyperglycemia (high blood sugar) or hypoglycemia (low blood sugar) related to the drugs occurred in fewer than one in 100 patients studied—but clinicians should consider the higher risk when treating diabetic patients with fluoroquinolones, especially moxifloxacin, and prescribe them cautiously, the study's authors concluded.

Increased use of these drugs, commonly used to treat such illnesses as [urinary tract infections](#) and community-acquired pneumonia, has raised concerns about rare but severe adverse effects, including tendon rupture and [heart arrhythmia](#). Previous studies have also indicated a relationship between fluoroquinolones and severe glucose-related abnormalities, known as [dysglycemia](#), which includes hyperglycemia or hypoglycemia. Severe blood sugar swings can lead to serious health problems, including irreversible brain damage or even death. In 2006, one drug from the fluoroquinolone class, gatifloxacin, was withdrawn from the U.S. market due to the risk of blood sugar abnormalities.

To assess the risk of blood sugar swings in diabetic patients using specific fluoroquinolones, a team of researchers, led by Mei-Shu Lai, MD, PhD, at National Taiwan University in Taipei, conducted a population-based cohort study of approximately 78,000 people with diabetes in Taiwan from January 2006 to November 2007.

Using the claims database for Taiwan's national

insurance program, the researchers analyzed data for diabetic outpatients who had received a new prescription for an antibiotic from one of three different classes of antibiotics: fluoroquinolones ([levofloxacin](#), ciprofloxacin, or moxifloxacin); second-generation cephalosporins (cefuroxime, cefaclor, or cefprozil); or macrolides (clarithromycin or azithromycin). The study's authors then looked for any emergency department visits or hospitalizations for dysglycemia among these patients within 30 days of the start of their antibiotic therapy.

Diabetics using oral fluoroquinolones faced greater risk of severe blood sugar swings than diabetic patients using antibiotics in other classes, the researchers found. The risks varied according to the specific fluoroquinolone the patients were using: The absolute risk, or incidence, of hyperglycemia per 1,000 people studied was 6.9 for moxifloxacin, 3.9 for levofloxacin, and 4.0 for ciprofloxacin. The absolute risk of hypoglycemia was 10.0 for moxifloxacin, 9.3 for levofloxacin, and 7.9 for ciprofloxacin.

(By comparison, among diabetic patients taking antibiotics in the macrolides class, the absolute risk of hyperglycemia was lower, at 1.6 per 1,000, and 2.1 per 1,000 among those taking antibiotics in the cephalosporin class; for hypoglycemia, the absolute risk per 1,000 was 3.7 for macrolides and 3.2 for cephalosporins, respectively.)

"Our results identified [moxifloxacin](#) as the drug associated with the highest risk of hypoglycemia, followed by levofloxacin and ciprofloxacin," the study's authors wrote. "Other antibiotics should be considered if dysglycemia is a concern, such as a beta lactam or macrolide," noted Dr. Lai.

Provided by Infectious Diseases Society of
America

APA citation: Study examines risk of severe blood sugar swings among diabetics taking fluoroquinolones
(2013, August 15) retrieved 21 November 2019 from [https://medicalxpress.com/news/2013-08-severe-
blood-sugar-diabetics-fluoroquinolones.html](https://medicalxpress.com/news/2013-08-severe-blood-sugar-diabetics-fluoroquinolones.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.