

# Combination of two imaging techniques allows new insights into brain function

August 26 2013

---

The ability to measure brain functions non-invasively is important both for clinical diagnoses and research in Neurology and Psychology. Two main imaging techniques are used: positron emission tomography (PET), which reveals metabolic processes in the brain; and activity of different brain regions is measured on the basis of the cells' oxygen consumption by magnetic resonance imaging (MRI). A direct comparison of PET and MRI measurements was previously difficult because each had to be performed in a separate machine.

Researchers from the Werner Siemens Imaging Center at the University of Tübingen under the direction of Professor Bernd J. Pichler in collaboration with the Department of Diagnostic and Interventional Radiology, University Hospital Tübingen, and the Tübingen Max Planck Institute for Intelligent Systems have now successfully combined both methods. The researchers are able to explore functional processes in the brain in detail and can better assess what course of action to take. These results were achieved by the use of a PET insert enabling complementary, simultaneous PET/MRI scans. It was developed and built at the University of Tübingen.

The researchers could identify in certain regions a mismatch between [glucose metabolism](#) related [brain activation](#) measured with PET and oxygenation related signals, measured with MRI. Furthermore information about [functional connectivity](#) in the brain could be derived from MRI and from dynamic PET data. These results help to further decipher the nature of brain function, and are ultimately useful for basic

research as well as clinical practice. The study, by lead author Dr. Hans Wehrl of Professor Bernd J. Pichler's research team has now been published in the journal *Nature Medicine*.

In PET imaging the distribution of a weakly radioactive substance is shown in cross sections of the body, enabling doctors to see many different metabolic and physiological functions at work. Functional MRI (fMRI) allows researchers to depict changes in blood oxygenation that are associated with brain function. This measurement of functional active brain regions is also important for the planning of brain surgeries, where particular care must be taken in certain areas. The ability to collect different kinds of data from different scans simultaneously represents a major step forward in the fields using these technologies.

**More information:** Wehrl, H. et al. Simultaneous PET-MRI reveals brain function in activated and resting state on metabolic, hemodynamic and multiple temporal scales, *Nature Medicine*. [DOI: 10.1038/nm.3290](https://doi.org/10.1038/nm.3290)

Provided by University of Tübingen

Citation: Combination of two imaging techniques allows new insights into brain function (2013, August 26) retrieved 20 September 2024 from <https://medicalxpress.com/news/2013-08-combination-imaging-techniques-insights-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.